

Anti-Histone H2B Rabbit Monoclonal Antibody

Catalog # ABO14087

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	O60814
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Histone H2B Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	85236
Other Names	Histone H2B type 1-K, H2B K, HIRA-interacting protein 1, H2BC12 (HGNC:13954)
Calculated MW	13890
Application Details	WB 1:5000-1:20000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:30 FC 1:50
Subcellular Localization	Nucleus. Chromosome.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HAD-8
Immunogen	A synthesized peptide derived from human Histone H2B
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	H2BC12 (HGNC:13954)
-------------	---------------------------------------


Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.

Images



Immunohistochemical analysis of paraffin-embedded human lung cancer, using Histone H2B Antibody.




Figure 1. Western blot analysis of Histone H2B using anti-Histone H2B antibody (M11068).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human A431 whole cell lysates,

Lane 2: human SIHA whole cell lysates,

Lane 3: rat brain tissue lysates,

Lane 4: rat liver tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Histone H2B antigen affinity purified monoclonal antibody (Catalog # M11068) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Histone H2B at approximately 17 kDa. The expected band size for Histone H2B is at 14 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.