10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-KMT6 / EZH2 Rabbit Monoclonal Antibody

Catalog # ABO14093

## Product Information

| | |
|---|---|
| **Application** | WB, IHC, IF, ICC, FC |
| **Primary Accession** | Q15910 |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Reactivity** | Rat, Human, Mouse |
| **Clonality** | Monoclonal |
| **Format** | Liquid |
| **Description** | Anti-KMT6 / EZH2 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. |


## Additional Information

| | |
|---|---|
| **Gene ID** | 2146 |
| **Other Names** | Histone-lysine N-methyltransferase EZH2, 2.1.1.356, ENX-1, Enhancer of zeste homolog 2, Lysine N-methyltransferase 6, EZH2 (HGNC:3527), KMT6 |
| **Calculated MW** | 85363 |
| **Application Details** | WB 1:500-1:1000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1 : 50 |
| **Subcellular Localization** | Nucleus. |
| **Tissue Specificity** | Expressed in many tissues. Overexpressed in numerous tumor types including carcinomas of the breast, colon, larynx, lymphoma and testis.. |
| **Source** | Eukaryota |
| **Contents** | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| **Clone Names** | Clone: IDF-5 |
| **Immunogen** | A synthesized peptide derived from human KMT6 / EZH2 |
| **Purification** | Affinity-chromatography |
| **Storage** | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |


## Protein Information

| | |
|---|---|
| **Name** | EZH2 ( HGNC:3527) |
| **Synonyms** | KMT6 |
| **Function** | Catalytic subunit of the PRC2/EED-EZH2 complex, a Polycomb group (PcG) complex that methylates 'Lys-9' (H3K9me) and 'Lys-27' (H3K27me) of histone H3, leading to transcriptional repression of the affected target gene (PubMed:14532106, PubMed:15225548, PubMed:15385962, PubMed:16618801, PubMed:16936726, PubMed:17344414, PubMed:22323599, PubMed:24474760, PubMed:26581166, PubMed:30026490, PubMed:30923826). Able to mono-, di- and trimethylate 'Lys-27' of histone H3 to form H3K27me1, H3K27me2 and H3K27me3, respectively (PubMed:15231737, PubMed:17210787, PubMed:18285464, PubMed:22323599, PubMed:30923826). Displays a preference for substrates with less methylation, loses activity when progressively more methyl groups are incorporated into H3K27, H3K27me0 > H3K27me1 > H3K27me2 (PubMed:22323599, PubMed:30923826). Compared to EZH1-containing complexes, it is more abundant in embryonic stem cells and plays a major role in forming H3K27me3, which is required for embryonic stem cell identity and proper differentiation (PubMed:19026781). The PRC2/EED-EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems (PubMed:16357870, PubMed:17200670). Genes repressed by the PRC2/EED- EZH2 complex include HOXC8, HOXA9, MYT1, CDKN2A and retinoic acid target genes (PubMed:16179254, PubMed:18086877, PubMed:20935635). EZH2 can also methylate non-histone proteins such as the transcription factor GATA4 and the nuclear receptor RORA (PubMed:23063525). Regulates the circadian clock via histone methylation at the promoter of the circadian genes (PubMed:16717091). Essential for the CRY1/2-mediated repression of the transcriptional activation of PER1/2 by the CLOCK- BMAL1 heterodimer; involved in the di and trimethylation of 'Lys-27' of histone H3 on PER1/2 promoters which is necessary for the CRY1/2 proteins to inhibit transcription (By similarity). |
| **Cellular Location** | Nucleus. Note=Localizes to the inactive X chromosome in trophoblast stem cells. {ECO:0000250\|UniProtKB:Q61188} |
| **Tissue Location** | In the ovary, expressed in primordial follicles and oocytes and also in external follicle cells (at protein level) (PubMed:31451685). Expressed in many tissues (PubMed:14532106) Overexpressed in numerous tumor types including carcinomas of the breast, colon, larynx, lymphoma and testis (PubMed:14532106) |


## Images




Western blot analysis of KMT6 / EZH2 expression in HEK293 cell lysate.



Immunohistochemical analysis of paraffin-embedded human tonsil, using KMT6 / EZH2 Antibody.




Immunofluorescent analysis of Hela cells, using KMT6 / EZH2 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.