

Anti-AMPK gamma 1 Rabbit Monoclonal Antibody

Catalog # ABO14125

Product Information

Application	WB, IP, FC
Primary Accession	P54619
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-AMPK gamma 1 Rabbit Monoclonal Antibody . Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	5571
Other Names	5'-AMP-activated protein kinase subunit gamma-1, AMPK gamma1, AMPK subunit gamma-1, AMPKg, PRKAG1
Calculated MW	37579
Application Details	WB 1:1000-1:2000 IP 1:20 FC 1:20
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: GDE-16
Immunogen	A synthesized peptide derived from human AMPK gamma 1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	PRKAG1
Function	AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism (PubMed: 21680840 , PubMed: 24563466). In response to

reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation (PubMed:[21680840](#), PubMed:[24563466](#)). AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators (PubMed:[21680840](#), PubMed:[24563466](#)). Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin (PubMed:[21680840](#), PubMed:[24563466](#)). Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits (PubMed:[21680840](#), PubMed:[24563466](#)). ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit (PubMed:[21680840](#), PubMed:[24563466](#)). ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive (PubMed:[21680840](#), PubMed:[24563466](#)).

Images



Western blot analysis of AMPK gamma 1 expression in HEK293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.