

Anti-14-3-3 gamma YWHAG Rabbit Monoclonal Antibody

Catalog # ABO14128

Product Information

Application	WB, FC
Primary Accession	P61981
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-14-3-3 gamma YWHAG Rabbit Monoclonal Antibody . Tested in WB, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	7532
Other Names	14-3-3 protein gamma, Protein kinase C inhibitor protein 1, KCIP-1, 14-3-3 protein gamma, N-terminally processed, YWHAG (HGNC:12852)
Calculated MW	28303
Application Details	WB 1:1000-1:2000 FC 1:50
Subcellular Localization	Cytoplasm.
Tissue Specificity	Highly expressed in brain, skeletal muscle, and heart..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HAH-25
Immunogen	A synthesized peptide derived from human 14-3-3 gamma
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	YWHAG (HGNC:12852)
-------------	--------------------------------------

Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:15696159, PubMed:16511572, PubMed:36732624). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:15696159, PubMed:16511572, PubMed:36732624). Binding generally results in the modulation of the activity of the binding partner (PubMed:16511572). Promotes inactivation of WDR24 component of the GATOR2 complex by binding to phosphorylated WDR24 (PubMed:36732624). Participates in the positive regulation of NMDA glutamate receptor activity by promoting the L- glutamate secretion through interaction with BEST1 (PubMed:29121962). Reduces keratinocyte intercellular adhesion, via interacting with PKP1 and sequestering it in the cytoplasm, thereby reducing its incorporation into desmosomes (PubMed:29678907). Plays a role in mitochondrial protein catabolic process (also named MALM) that promotes the degradation of damaged proteins inside mitochondria (PubMed:22532927).
Cellular Location	Cytoplasm, cytosol. Mitochondrion matrix. Note=Translocates to the mitochondrial matrix following induction of MALM (mitochondrial protein catabolic process).
Tissue Location	Highly expressed in brain, skeletal muscle, and heart.

Images



Western blot analysis of 14-3-3 gamma expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.