

Anti-14-3-3 sigma SFN Rabbit Monoclonal Antibody

Catalog # ABO14129

Product Information

Application	WB, IHC
Primary Accession	P31947
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-14-3-3 sigma SFN Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Rat.


Additional Information

Gene ID	2810
Other Names	14-3-3 protein sigma, Epithelial cell marker protein 1, Stratifin, SFN, HME1 {ECO:0000303 PubMed:1390337}
Calculated MW	27774
Application Details	WB 1:1000-1:2000 IHC 1:50-1:200
Subcellular Localization	Cytoplasm. Nucleus. Secreted. May be secreted by a non-classical secretory pathway.
Tissue Specificity	Present mainly in tissues enriched in stratified squamous keratinizing epithelium.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HBB-19
Immunogen	A synthesized peptide derived from human 14-3-3 sigma
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	SFN
Synonyms	HME1 {ECO:0000303 PubMed:1390337}
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Binding generally results in the modulation of the activity of the binding partner (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Promotes cytosolic retention of GBP1 GTPase by binding to phosphorylated GBP1, thereby inhibiting the innate immune response (PubMed:37797010). Also acts as a TP53/p53-regulated inhibitor of G2/M progression (PubMed:9659898). When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity). Acts to maintain desmosome cell junction adhesion in epithelial cells via interacting with and sequestering PKP3 to the cytoplasm, thereby restricting its translocation to existing desmosome structures and therefore maintaining desmosome protein homeostasis (PubMed:24124604). Also acts to facilitate PKP3 exchange at desmosome plaques, thereby maintaining keratinocyte intercellular adhesion (PubMed:29678907). May also regulate MDM2 autoubiquitination and degradation and thereby activate p53/TP53 (PubMed:18382127).
Cellular Location	Cytoplasm. Nucleus {ECO:0000250 UniProtKB:O70456} Secreted. Note=May be secreted by a non- classical secretory pathway.
Tissue Location	Present mainly in tissues enriched in stratified squamous keratinizing epithelium.

Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of 14-3-3 sigma expression in A431 cell lysate.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.