

Anti-CaMKII alpha CAMK2A Rabbit Monoclonal Antibody

Catalog # ABO14133

Product Information

Application	WB, IP, FC
Primary Accession	Q9UQM7
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CaMKII alpha CAMK2A Rabbit Monoclonal Antibody . Tested in WB, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	815
Other Names	Calcium/calmodulin-dependent protein kinase type II subunit alpha, CaM kinase II subunit alpha, CaMK-II subunit alpha, 2.7.11.17, CAMK2A, CAMKA, KIAA0968
Calculated MW	54088
Application Details	WB 1:1000-1:5000 IP 1:50 FC 1:100
Subcellular Localization	Cell junction, synapse, presynaptic cell membrane. Cell junction, synapse. Postsynaptic lipid rafts..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAFF-3
Immunogen	A synthesized peptide derived from human CaMKII alpha
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CAMK2A
-------------	--------

Synonyms

CAMKA, KIAA0968


Function

Calcium/calmodulin-dependent protein kinase that functions autonomously after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in various processes, such as synaptic plasticity, neurotransmitter release and long-term potentiation (PubMed:[14722083](#)). Member of the NMDAR signaling complex in excitatory synapses, it regulates NMDAR-dependent potentiation of the AMPAR and therefore excitatory synaptic transmission (By similarity). Regulates dendritic spine development (PubMed:[28130356](#)). Also regulates the migration of developing neurons (PubMed:[29100089](#)). Phosphorylates the transcription factor FOXO3 to activate its transcriptional activity (PubMed:[23805378](#)). Phosphorylates the transcription factor ETS1 in response to calcium signaling, thereby decreasing ETS1 affinity for DNA (By similarity). In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK- STAT signaling pathway (PubMed:[11972023](#)). In response to interferon- beta (IFN-beta) stimulation, stimulates the JAK-STAT signaling pathway (PubMed:[35568036](#)). In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of PSAT1, inhibiting ferroptosis by promoting GPX4 hydroxylation and stability (PubMed:[40281343](#)). Acts as a negative regulator of 2-arachidonoylglycerol (2-AG)-mediated synaptic signaling via modulation of DAGLA activity (By similarity).

Cellular Location

Synapse {ECO:0000250|UniProtKB:P11275}. Postsynaptic density {ECO:0000250|UniProtKB:P11275}. Cell projection, dendritic spine. Cell projection, dendrite. Note=Postsynaptic lipid rafts {ECO:0000250|UniProtKB:P11275}

Images



Western blot analysis of CaMKII alpha expression in SH-SY5Y cell lysate.




Figure 1. Western blot analysis of CAMK2A using anti-CAMK2A antibody (M03241). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat Hippocampus tissue lysates, Lane 2: rat brain tissue lysates, Lane 3: rat C6 whole cell lysates, Lane 4: mouse brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CAMK2A antigen affinity purified monoclonal

antibody (Catalog # M03241) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CAMK2A at approximately 54 kDa. The expected band size for CAMK2A is at 54 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.