

Anti-Cytokeratin 1 Rabbit Monoclonal Antibody

Catalog # ABO14156

Product Information

Application	WB, IHC
Primary Accession	P04264
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Cytokeratin 1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	3848
Other Names	Keratin, type II cytoskeletal 1, 67 kDa cytokeratin, Cytokeratin-1, CK-1, Hair alpha protein, Keratin-1, K1, Type-II keratin Kb1, KRT1, KRTA
Calculated MW	66039
Application Details	WB 1:5000-1:20000 IHC 1:50-1:200
Subcellular Localization	Cell membrane. Located on plasma membrane of neuroblastoma NMB7 cells.
Tissue Specificity	The source of this protein is neonatal foreskin. The 67-kDa type II keratins are expressed in terminally differentiating epidermis.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: GBH-11
Immunogen	A synthesized peptide derived from human Cytokeratin 1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	KRT1 (HGNC:6412)
Synonyms	KRTA
Function	Structural component of intermediate filaments in suprabasal keratinocytes of stratified epithelia. Forms heteropolymers with a type I keratin, assembling into keratin intermediate filament networks that provide mechanical strength and structural stability to differentiating epidermal cells (PubMed: 1381288). May regulate the activity of kinases such as PKC and SRC by interacting with integrin beta-1 (ITB1) and the receptor of activated protein C kinase 1 (RACK1) (PubMed: 17956333 , PubMed: 21544310). In complex with C1QBP, acts as a high-affinity receptor for kininogen-1 (HMWK) (PubMed: 21544310).
Cellular Location	Cell membrane. Cytoplasm
Tissue Location	The source of this protein is neonatal foreskin. The 67-kDa type II keratins are expressed in terminally differentiating epidermis


Images



All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.



Western blot analysis of Cytokeratin 1 expression in Human skin lysate.



Immunohistochemical analysis of paraffin-embedded rat skin, using Cytokeratin 1 Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.