

Anti-TRK fused gene TFG Rabbit Monoclonal Antibody

Catalog # ABO14194

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q92734
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TRK fused gene TFG Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	10342
Other Names	Protein TFG, TRK-fused gene protein, TFG
Calculated MW	43448
Application Details	WB 1:5000-1:20000 IHC 1:50-1:200 ICC/IF 1:100-1:500 IP 1:50
Tissue Specificity	Ubiquitous.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HFB-20
Immunogen	A synthesized peptide derived from human TRK fused gene
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	TFG
Function	Plays a role in the normal dynamic function of the endoplasmic reticulum

(ER) and its associated microtubules (PubMed:[23479643](#), PubMed:[27813252](#)). Required for secretory cargo traffic from the endoplasmic reticulum to the Golgi apparatus (PubMed:[21478858](#)).


Cellular Location

Endoplasmic reticulum. Note=Localizes to endoplasmic reticulum exit site (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:21478858, PubMed:27813252).

Tissue Location

Ubiquitous.

Images



Western blot analysis of TRK fused gene expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.