

Anti-Alpha Synuclein SNCA Rabbit Monoclonal Antibody

Catalog # ABO14203

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P37840
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Alpha Synuclein SNCA Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.


Additional Information

Gene ID	6622
Other Names	Alpha-synuclein, Non-A beta component of AD amyloid, Non-A4 component of amyloid precursor, NACP, SNCA, NACP, PARK1
Calculated MW	14460
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Subcellular Localization	Cytoplasm, cytosol. Membrane. Nucleus. Cell junction, synapse. Secreted. Membrane-bound in dopaminergic neurons..
Tissue Specificity	Expressed principally in brain but is also expressed in low concentrations in all tissues examined except in liver. Concentrated in presynaptic nerve terminals.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AGA-19
Immunogen	A synthesized peptide derived from human Alpha Synuclein
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information



Name	SNCA
Synonyms	NACP, PARK1
Function	Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed: 20798282 , PubMed: 26442590 , PubMed: 28288128 , PubMed: 30404828). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed: 28288128 , PubMed: 30404828). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed: 30404828). Also acts as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5 (PubMed: 20798282). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed: 20798282). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed: 26442590).
Cellular Location	Cytoplasm. Membrane Nucleus Synapse. Secreted. Cell projection, axon {ECO:0000250 UniProtKB:O55042}. Note=Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). {ECO:0000250 UniProtKB:O55042, ECO:0000269 PubMed:15282274}
Tissue Location	Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain

Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.


Western blot analysis of Alpha Synuclein expression in Human fetal brain lysate.




Immunohistochemical analysis of paraffin-embedded mouse testis, using Alpha Synuclein Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis using the Antibody at 1:150 dilution.



Immunofluorescent analysis using the Antibody at 1:500 dilution.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.