

Anti-Integrin alpha V ITGAV Rabbit Monoclonal Antibody

Catalog # ABO14232

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	P06756
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Integrin alpha V ITGAV Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	3685
Other Names	Integrin alpha-V, Vitronectin receptor {ECO:0000312 HGNC:HGNC:6150}, Vitronectin receptor subunit alpha, CD51, Integrin alpha-V heavy chain, Integrin alpha-V light chain, ITGAV (HGNC:6150)
Calculated MW	116038
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:40
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABGC-9
Immunogen	A synthesized peptide derived from human Integrin alpha V
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ITGAV (HGNC:6150)
-------------	-------------------------------------

Function

The alpha-V (ITGAV) integrins are receptors for vitronectin, cytotactin, fibronectin, fibrinogen, laminin, matrix metalloproteinase- 2, osteopontin, osteomodulin, prothrombin, thrombospondin and vWF. They recognize the sequence R-G-D in a wide array of ligands. ITGAV:ITGB3 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1- dependent fractalkine signaling (PubMed:[23125415](#)). ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:[20682778](#)). ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling (PubMed:[18441324](#)). ITGAV:ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling (PubMed:[28302677](#)). ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:[19578119](#)). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:[28873464](#)). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (PubMed:[29030430](#)). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:[18635536](#), PubMed:[25398877](#)). ITGAV:ITGB3 and ITGAV:ITGB6 act as receptors for fibrillin-1 (FBN1) and mediate R-G-D-dependent cell adhesion to FBN1 (PubMed:[12807887](#), PubMed:[17158881](#)). Integrin alpha-V/beta-6 or alpha- V/beta-8 (ITGAV:ITGB6 or ITGAV:ITGB8) mediates R-G-D-dependent release of transforming growth factor beta-1 (TGF-beta-1) from regulatory Latency-associated peptide (LAP), thereby playing a key role in TGF- beta-1 activation (PubMed:[15184403](#), PubMed:[22278742](#), PubMed:[28117447](#)). ITGAV:ITGB3 acts as a receptor for CD40LG (PubMed:[31331973](#)). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (PubMed:[10640428](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell junction, focal adhesion

Images




Figure 1. Western blot analysis of ITGAV using anti-ITGAV antibody (M01561).


Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human A549 whole cell lysates,
Lane 2: human A431 whole cell lysates,
Lane 3: human 293T whole cell lysates,
Lane 4: human MCF-7 whole cell lysates,
Lane 5: rat spleen tissue lysates,
Lane 6: rat brain tissue lysates,
Lane 7: mouse spleen tissue lysates,
Lane 8: mouse brain tissue lysates.


After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ITGAV antigen affinity purified monoclonal antibody (Catalog # M01561) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ITGAV at

approximately 130, 140, 150 kDa. The expected band size for ITGAV is at 116 kDa.


Immunofluorescent analysis using the Antibody at 1:50 dilution.




Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis using the Antibody at 1:150 dilution.



Immunofluorescent analysis using the Antibody at 1:500 dilution.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.