

Anti-Lipoprotein lipase LPL Rabbit Monoclonal Antibody

Catalog # ABO14261

Product Information

Application	WB, IHC
Primary Accession	P06858
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Lipoprotein lipase LPL Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Additional Information


Gene ID	4023
Other Names	Lipoprotein lipase, LPL, 3.1.1.34, Phospholipase A1, 3.1.1.32, LPL, LIPD
Calculated MW	53162
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200
Subcellular Localization	Cell membrane ; Lipid-anchor, GPI-anchor. Secreted. Locates to the plasma membrane of microvilli of hepatocytes with triacyl- glycerol-rich lipoproteins (TRL). Some of the bound LPL is then internalized and located inside non-coated endocytic vesicles (By similarity)..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ABDE-12
Immunogen	A synthesized peptide derived from human Lipoprotein lipase
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	LPL
-------------	-----

Synonyms	LIPD
Function	Key enzyme in triglyceride metabolism. Catalyzes the hydrolysis of triglycerides from circulating chylomicrons and very low density lipoproteins (VLDL), and thereby plays an important role in lipid clearance from the blood stream, lipid utilization and storage (PubMed: 11342582 , PubMed: 27578112 , PubMed: 8675619). Although it has both phospholipase and triglyceride lipase activities it is primarily a triglyceride lipase with low but detectable phospholipase activity (PubMed: 12032167 , PubMed: 7592706). Mediates margination of triglyceride-rich lipoprotein particles in capillaries (PubMed: 24726386). Recruited to its site of action on the luminal surface of vascular endothelium by binding to GPIHBP1 and cell surface heparan sulfate proteoglycans (PubMed: 11342582 , PubMed: 27811232).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P11151}; Peripheral membrane protein {ECO:0000250 UniProtKB:P11151}; Extracellular side {ECO:0000250 UniProtKB:P11151}. Secreted. Secreted, extracellular space, extracellular matrix. Note=Newly synthesized LPL binds to cell surface heparan proteoglycans and is then released by heparanase Subsequently, it becomes attached to heparan proteoglycan on endothelial cells (PubMed:27811232). Locates to the plasma membrane of microvilli of hepatocytes with triglyceride-rich lipoproteins (TRL) Some of the bound LPL is then internalized and located inside non- coated endocytic vesicles (By similarity) {ECO:0000250 UniProtKB:P11151, ECO:0000269 PubMed:27811232}
Tissue Location	Detected in blood plasma (PubMed:11893776, PubMed:12641539, PubMed:2340307). Detected in milk (at protein level) (PubMed:2340307).


Images




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of Lipoprotein lipase expression in Human fetal liver lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.