

Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody

Catalog # ABO14264

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P19544
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.


Additional Information

Gene ID	7490
Other Names	Wilms tumor protein, WT33, WT1
Calculated MW	49188
Application Details	WB 1:500-1:1000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Subcellular Localization	Nucleus. Nucleus, nucleolus. Cytoplasm. Shuttles between nucleus and cytoplasm..
Tissue Specificity	Expressed in the kidney and a subset of hematopoietic cells.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: CEH-23
Immunogen	A synthesized peptide derived from human Wilms Tumor Protein
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	WT1
Function	Transcription factor that plays an important role in cellular development and cell survival (PubMed: 7862533). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed: 17716689 , PubMed: 25258363 , PubMed: 7862533). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific; isoforms lacking the KTS motif may act as transcription factors (PubMed: 15520190). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed: 16934801). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed: 19123921).
Cellular Location	Nucleus. Nucleus, nucleolus. Cytoplasm. Note=Isoforms lacking the KTS motif have a diffuse nuclear location (PubMed:15520190). Shuttles between nucleus and cytoplasm. {ECO:0000250, ECO:0000269 PubMed:15520190} [Isoform 4]: Nucleus, nucleoplasm
Tissue Location	Expressed in the kidney and a subset of hematopoietic cells

Images





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Western blot analysis of WT1 expression in K562 cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse kidney, using Wilms Tumor Protein Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.