

Anti-Neutrophil Elastase ELANE Rabbit Monoclonal Antibody

Catalog # ABO14267

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P08246
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Neutrophil Elastase ELANE Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.


Additional Information

Gene ID	1991
Other Names	Neutrophil elastase, 3.4.21.37, Bone marrow serine protease, Elastase-2, Human leukocyte elastase, HLE, Medullasin, PMN elastase, ELANE, ELA2
Calculated MW	28518
Application Details	WB 1:500-1:2000 IHC 1:100-1:500 ICC/IF 1:50-1:200 FC 1:30
Tissue Specificity	Bone marrow cells.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AOCI-5
Immunogen	A synthesized peptide derived from human Neutrophil Elastase
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	ELANE
Synonyms	ELA2
Function	Serine protease that modifies the functions of natural killer cells, monocytes and granulocytes (PubMed: 15140022). Shows microbicidal activity in neutrophils by mediating cleavage and degradation of bacterial proteins (PubMed: 10947984 , PubMed: 25161283). Capable of killing E.coli but not S.aureus in vitro; digests outer membrane protein A (ompA) in E.coli and K.pneumoniae (PubMed: 10947984). Involved for the formation of neutrophil extracellular traps (NETs): translocates into the nucleus downstream of MPO and mediates degradation of histone H4, thereby participating to chromatin decondensation (PubMed: 20974816 , PubMed: 25066128 , PubMed: 28574339). NETs are mainly composed of DNA fibers and are released by neutrophils to trap pathogens during inflammation (PubMed: 20974816 , PubMed: 25066128 , PubMed: 28574339). Inhibits C5a-dependent neutrophil enzyme release and chemotaxis (PubMed: 15140022). Promotes cleavage of GSDMB, thereby inhibiting pyroptosis (PubMed: 36899106). Promotes blood coagulation (PubMed: 20676107). Through the activation of the platelet fibrinogen receptor integrin alpha-IIb/beta-3, potentiates platelet aggregation induced by a threshold concentration of cathepsin G (CTSG) (PubMed: 25211214 , PubMed: 9111081). Cleaves and thus inactivates tissue factor pathway inhibitor (TFPI) (PubMed: 20676107 , PubMed: 25211214).
Cellular Location	Cytoplasmic vesicle, phagosome. Lysosome Cytolytic granule. Nucleus. Note=Localizes to azurophil granules, a type of primary lysosome found in neutrophils and monocytes that contains antimicrobial proteins (PubMed:20974816, PubMed:25066128) Translocates from azurophil granules to the nucleus downstream of MPO during formation of neutrophil extracellular traps (NETs), and is present in released NETs (PubMed:20974816, PubMed:25066128, PubMed:32434865). Localized in phagolysosomes following ingestion of E.coli by neutrophils (PubMed:10947984).
Tissue Location	Bone marrow cells. Neutrophil (PubMed:10947984).

Images



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.

Figure 1. Western blot analysis of Neutrophil Elastase using anti-Neutrophil Elastase antibody (M01607). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.
Lane 1: human HL-60 whole cell lysates,
Lane 2: human THP-1 whole cell lysates.



After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Neutrophil Elastase antigen affinity purified monoclonal antibody (Catalog # M01607) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Neutrophil Elastase at approximately 29 kDa. The expected band size for Neutrophil Elastase is at 29 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.