10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Myosin light chain kinase MYLK Rabbit Monoclonal Antibody

Catalog # ABO14308

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q15746
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Myosin light chain kinase MYLK Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4638
Other Names	Myosin light chain kinase, smooth muscle, MLCK, smMLCK, 2.7.11.18, Kinase-related protein, KRP, Telokin, Myosin light chain kinase, smooth muscle, deglutamylated form, MYLK (HGNC:7590), MLCK, MLCK1, MYLK1
Calculated MW	210715
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Subcellular Localization	Cytoplasm. Cell projection, lamellipodium. Cleavage furrow. Cytoplasm, cytoskeleton. Localized to stress fibers during interphase and to the cleavage furrow during mitosis.
Tissue Specificity	Smooth muscle and non-muscle isozymes are expressed in a wide variety of adult and fetal tissues and in cultured endothelium with qualitative expression appearing to be neither tissue- nor development-specific. Non-muscle isoform 2 is the dominant splice variant expressed in various tissues. Telokin has been found in a wide variety of adult and fetal tissues. Accumulates in well differentiated enterocytes of the intestinal epithelium in response to tumor necrosis factor (TNF)..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.


Clone Names	Clone: G-13
Immunogen	A synthesized peptide derived from human Myosin light chain kinase
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	MYLK (HGNC:7590)
Synonyms	MLCK, MLCK1, MYLK1
Function	Calcium/calmodulin-dependent myosin light chain kinase implicated in smooth muscle contraction via phosphorylation of myosin light chains (MLC) (PubMed:11113114). Also regulates actin-myosin interaction through a non-kinase activity (PubMed:11113114). Phosphorylates PTK2B/PYK2 and myosin light-chains (PubMed:11113114, PubMed:18587400). Involved in the inflammatory response (e.g. apoptosis, vascular permeability, leukocyte diapedesis), cell motility and morphology, airway hyperreactivity and other activities relevant to asthma (PubMed:20139351). Required for tonic airway smooth muscle contraction that is necessary for physiological and asthmatic airway resistance (By similarity). Necessary for gastrointestinal motility (By similarity). Implicated in the regulation of endothelial as well as vascular permeability, probably via the regulation of cytoskeletal rearrangements (By similarity). In the nervous system it has been shown to control the growth initiation of astrocytic processes in culture and to participate in transmitter release at synapses formed between cultured sympathetic ganglion cells (PubMed:20375339). Critical participant in signaling sequences that result in fibroblast apoptosis (PubMed:20453870). Plays a role in the regulation of epithelial cell survival (PubMed:16723733). Required for epithelial wound healing, especially during actomyosin ring contraction during purse-string wound closure (PubMed:15825080). Mediates RhoA-dependent membrane blebbing (PubMed:20181817). Triggers TRPC5 channel activity in a calcium-dependent signaling, by inducing its subcellular localization at the plasma membrane (PubMed:16284075). Promotes cell migration (including tumor cells) and tumor metastasis (PubMed:18710790, PubMed:19826488). PTK2B/PYK2 activation by phosphorylation mediates ITGB2 activation and is thus essential to trigger neutrophil transmigration during acute lung injury (ALI) (PubMed:18587400). May regulate optic nerve head astrocyte migration (PubMed:20375339). Probably involved in mitotic cytoskeletal regulation (PubMed:15020676). Regulates tight junction probably by modulating ZO-1 exchange in the perijunctional actomyosin ring (By similarity). Mediates burn-induced microvascular barrier injury; triggers endothelial contraction in the development of microvascular hyperpermeability by phosphorylating MLC (By similarity). Essential for intestinal barrier dysfunction (By similarity). Mediates Giardia spp.-mediated reduced epithelial barrier function during giardiasis intestinal infection via reorganization of cytoskeletal F- actin and tight junctional ZO-1 (By similarity). Necessary for hypotonicity-induced Ca(2+) entry and subsequent activation of volume- sensitive organic osmolyte/anion channels (VSOAC) in cervical cancer cells (PubMed:11976941). Responsible for high proliferative ability of breast cancer cells through anti-apoptosis (PubMed:20453870).
Cellular Location	Cytoplasm. Cell projection, lamellipodium. Cleavage furrow. Cytoplasm, cytoskeleton, stress fiber. Note=Localized to stress fibers during interphase and to the cleavage furrow during mitosis

Tissue Location	Smooth muscle and non-muscle isozymes are expressed in a wide variety of adult and fetal tissues and in cultured endothelium with qualitative expression appearing to be neither tissue- nor development-specific. Non-muscle isoform 2 is the dominant splice variant expressed in various tissues. Telokin has been found in a wide variety of adult and fetal tissues. Accumulates in well differentiated enterocytes of the intestinal epithelium in response to tumor necrosis factor (TNF).

Images



Western blot analysis of myosin light chain kinase expression in HUVEC cell lysate.



Immunohistochemical analysis of paraffin-embedded human colon, using Myosin light chain kinase Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.