

Anti-Interferon Receptor alpha IFNAR1 Rabbit Monoclonal Antibody

Catalog # ABO14310

Product Information

Application	WB
Primary Accession	P17181
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Interferon Receptor alpha IFNAR1 Rabbit Monoclonal Antibody . Tested in WB application. This antibody reacts with Human.

Additional Information

Gene ID	3454
Other Names	Interferon alpha/beta receptor 1, IFN-R-1, IFN-alpha/beta receptor 1, Cytokine receptor class-II member 1, Cytokine receptor family 2 member 1, CRF2-1, Type I interferon receptor 1, IFNAR1, IFNAR
Calculated MW	63525
Application Details	WB 1:500-1:2000
Subcellular Localization	Isoform 1: Cell membrane ; Single- pass type I membrane protein. Late endosome. Lysosome. Interferon binding triggers internalization of the receptor from the cell membrane into endosomes and then into lysosomes..
Tissue Specificity	IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1, isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266B1. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN-alpha resistant myeloma cell line U266R..
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AF-9
Immunogen	A synthesized peptide derived from human Interferon Receptor alpha
Purification	Affinity-chromatography

Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
----------------	--

Protein Information

Name	IFNAR1
-------------	--------


Synonyms	IFNAR
-----------------	-------

Function	Together with IFNAR2, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed: 10049744 , PubMed: 14532120 , PubMed: 15337770 , PubMed: 2153461 , PubMed: 21854986 , PubMed: 24075985 , PubMed: 31270247 , PubMed: 33252644 , PubMed: 35442418 , PubMed: 7813427). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response (PubMed: 10049744 , PubMed: 21854986 , PubMed: 7665574). Mechanistically, type I interferon- binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed: 21854986 , PubMed: 32972995 , PubMed: 7665574 , PubMed: 7813427). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription factors (PubMed: 21854986 , PubMed: 32972995 , PubMed: 7526154 , PubMed: 7665574 , PubMed: 7813427). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed: 19561067 , PubMed: 21854986 , PubMed: 32972995 , PubMed: 7665574 , PubMed: 7813427 , PubMed: 9121453). Can also act independently of IFNAR2: form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).
-----------------	--

Cellular Location	[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Late endosome. Lysosome. Note=Interferon binding triggers internalization of the receptor from the cell membrane into endosomes and then into lysosomes.
--------------------------	---

Tissue Location	IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1, isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266B1. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN- alpha resistant myeloma cell line U266R.
------------------------	---

Images



Western blot analysis of Interferon Receptor alpha expression in SH-SY5Y cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.