

Anti-Histone H3 (mono methyl R17) HIST1H3A Rabbit Monoclonal Antibody

Catalog # ABO14317

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P68431
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Histone H3 (mono methyl R17) HIST1H3A Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Gene ID	8350;8351;8352;8353;8354;8355;8356;8357;8358;8968
Other Names	Histone H3.1, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, H3C1 (HGNC:4766), H3FA, HIST1H3A
Calculated MW	15404
Application Details	WB 1:100-1:2000 IHC 1:100-1:250 ICC/IF 1:100-1:250
Subcellular Localization	Nucleus. Chromosome.
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BDI-8
Immunogen	A synthesized peptide derived from human Histone H3 (mono methyl R17)
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Name	H3C1 (HGNC:4766)
Synonyms	H3FA, HIST1H3A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.


Images



Western blot analysis of Histone H3 (mono methyl R17) expression in HeLa cell lysate.




Immunohistochemical analysis of paraffin-embedded mouse colon, using Histone H3 (mono methyl R17) Antibody.




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.