

Anti-HIP1 Monoclonal Antibody

Catalog # ABO14354

Product Information

Application	WB, IF, ICC, IP
Primary Accession	O00291
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-HIP1 Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	3092
Other Names	Huntingtin-interacting protein 1, HIP-1, Huntingtin-interacting protein I, HIP-I, HIP1
Calculated MW	116221
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:30
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: AFG-8 A synthesized peptide derived from human HIP1 Plays a role in clathrin-mediated endocytosis and trafficking. Involved in regulating AMPA receptor trafficking in the central nervous system in an NMDA-dependent manner. Enhances androgen receptor (AR) -mediated transcription.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	HIP1
-------------	------

Function

Plays a role in clathrin-mediated endocytosis and trafficking (PubMed:[11532990](#), PubMed:[11577110](#), PubMed:[11889126](#)). Involved in regulating AMPA receptor trafficking in the central nervous system in an NMDA-dependent manner (By similarity). Regulates presynaptic nerve terminal activity (By similarity). Enhances androgen receptor (AR)- mediated transcription (PubMed:[16027218](#)). May act as a proapoptotic protein that induces cell death by acting through the intrinsic apoptosis pathway (PubMed:[11007801](#)). Binds 3-phosphoinositides (via ENTH domain) (PubMed:[14732715](#)). May act through the ENTH domain to promote cell survival by stabilizing receptor tyrosine kinases following ligand-induced endocytosis (PubMed:[14732715](#)). May play a functional role in the cell filament networks (PubMed:[18790740](#)). May be required for differentiation, proliferation, and/or survival of somatic and germline progenitors (PubMed:[11007801](#), PubMed:[12163454](#)).

Cellular Location

Cytoplasm. Nucleus. Endomembrane system. Cytoplasmic vesicle, clathrin-coated vesicle membrane. Note=Shuttles between cytoplasm and nucleus. Nuclear translocation can be induced by AR

Tissue Location

Ubiquitously expressed with the highest level in brain. Expression is up-regulated in prostate and colon cancer

Images




Figure 1. Western blot analysis of HIP1 using anti-HIP1 antibody (M02242).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,
Lane 2: human MCF-7 whole cell lysates,
Lane 3: human PC-3 whole cell lysates,
Lane 4: rat brain tissue lysates,
Lane 5: rat PC-12 whole cell lysates,
Lane 6: mouse brain tissue lysates,
Lane 7: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HIP1 antigen affinity purified monoclonal antibody (Catalog # M02242) at 1:1000 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for HIP1 at approximately 116 kDa. The expected band size for HIP1 is at 116 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.