

Anti-ATM Monoclonal Antibody

Catalog # ABO14357

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q13315
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-ATM Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	472
Other Names	Serine-protein kinase ATM, 2.7.11.1, Ataxia telangiectasia mutated, A-T mutated, ATM
Calculated MW	350687
Application Details	WB 1:500-1:2000 IHC 1:50-1:100 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: BCF-1
Immunogen	A synthesized peptide derived from human ATM
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ATM
Function	Serine/threonine protein kinase which activates checkpoint signaling upon double strand breaks (DSBs), apoptosis and genotoxic stresses such as

ionizing ultraviolet A light (UVA), thereby acting as a DNA damage sensor (PubMed:[10550055](#), PubMed:[10839545](#), PubMed:[10910365](#), PubMed:[12556884](#), PubMed:[14871926](#), PubMed:[15064416](#), PubMed:[15448695](#), PubMed:[15456891](#), PubMed:[15790808](#), PubMed:[15916964](#), PubMed:[17923702](#), PubMed:[21757780](#), PubMed:[24534091](#), PubMed:[27941124](#), PubMed:[35076389](#), PubMed:[9733514](#)). Recognizes the substrate consensus sequence [ST]-Q (PubMed:[10550055](#), PubMed:[10839545](#), PubMed:[10910365](#), PubMed:[12556884](#), PubMed:[14871926](#), PubMed:[15448695](#), PubMed:[15456891](#), PubMed:[15916964](#), PubMed:[17923702](#), PubMed:[24534091](#), PubMed:[27941124](#), PubMed:[9733514](#)). Phosphorylates 'Ser-139' of histone variant H2AX at double strand breaks (DSBs), thereby regulating DNA damage response mechanism (By similarity). Also plays a role in pre-B cell allelic exclusion, a process leading to expression of a single immunoglobulin heavy chain allele to enforce clonality and monospecific recognition by the B-cell antigen receptor (BCR) expressed on individual B-lymphocytes. After the introduction of DNA breaks by the RAG complex on one immunoglobulin allele, acts by mediating a repositioning of the second allele to pericentromeric heterochromatin, preventing accessibility to the RAG complex and recombination of the second allele. Also involved in signal transduction and cell cycle control. May function as a tumor suppressor. Necessary for activation of ABL1 and SAPK. Phosphorylates DYRK2, CHEK2, p53/TP53, FBXW7, FANCD2, NFKBIA, BRCA1, CREBBP/CBP, RBBP8/CTIP, FBXO46, MRE11, nibrin (NBN), RAD50, RAD17, PELI1, TERF1, UFL1, RAD9, UBQLN4, UCHL3 and DCLRE1C (PubMed:[10550055](#), PubMed:[10766245](#), PubMed:[10802669](#), PubMed:[10839545](#), PubMed:[10910365](#), PubMed:[10973490](#), PubMed:[11375976](#), PubMed:[12086603](#), PubMed:[15456891](#), PubMed:[19965871](#), PubMed:[21757780](#), PubMed:[24534091](#), PubMed:[26240375](#), PubMed:[26774286](#), PubMed:[27941124](#), PubMed:[30171069](#), PubMed:[30612738](#), PubMed:[30886146](#), PubMed:[30952868](#), PubMed:[38128537](#), PubMed:[9733515](#), PubMed:[9843217](#)). May play a role in vesicle and/or protein transport. Could play a role in T-cell development, gonad and neurological function. Plays a role in replication-dependent histone mRNA degradation. Binds DNA ends. Phosphorylation of DYRK2 in nucleus in response to genotoxic stress prevents its MDM2-mediated ubiquitination and subsequent proteasome degradation (PubMed:[19965871](#)). Phosphorylates ATF2 which stimulates its function in DNA damage response (PubMed:[15916964](#)). Phosphorylates ERCC6 which is essential for its chromatin remodeling activity at DNA double-strand breaks (PubMed:[29203878](#)). Phosphorylates TTC5/STRAP at 'Ser-203' in the cytoplasm in response to DNA damage, which promotes TTC5/STRAP nuclear localization (PubMed:[15448695](#)). Also involved in pexophagy by mediating phosphorylation of PEX5: translocated to peroxisomes in response to reactive oxygen species (ROS), and catalyzes phosphorylation of PEX5, promoting PEX5 ubiquitination and induction of pexophagy (PubMed:[26344566](#)).


Cellular Location

Nucleus. Cytoplasmic vesicle. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q62388}. Peroxisome matrix. Note=Primarily nuclear (PubMed:9050866, PubMed:9150358). Found also in endocytic vesicles in association with beta-adaptin (PubMed:9707615). Translocated to peroxisomes in response to reactive oxygen species (ROS) by PEX5 (PubMed:26344566)


Tissue Location

Found in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain, heart, spleen, thymus, testis, ovary, small intestine, colon and leukocytes

Images



Immunohistochemical analysis of paraffin-embedded human breast cancer, using ATM Antibody.



Immunofluorescent analysis of HeLa cells, using ATM Antibody.




Figure 1. Western blot analysis of ATM using anti-ATM antibody (M00014).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human 293T whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ATM antigen affinity purified monoclonal antibody (Catalog # M00014) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for ATM at approximately 351 kDa. The expected band size for ATM is at 351 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.