

Anti-RUNX2 Monoclonal Antibody

Catalog # ABO14378

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	Q13950
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-RUNX2 Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	860
Other Names	Runt-related transcription factor 2, Acute myeloid leukemia 3 protein, Core-binding factor subunit alpha-1, CBF-alpha-1, Oncogene AML-3, Osteoblast-specific transcription factor 2, OSF-2, Polyomavirus enhancer-binding protein 2 alpha A subunit, PEA2-alpha A, PEBP2-alpha A, SL3-3 enhancer factor 1 alpha A subunit, SL3/AKV core-binding factor alpha A subunit, RUNX2, AML3, CBFA1, OSF2, PEBP2A
Calculated MW	56648
Application Details	WB 1:500-1:2000 IHC 1:100-1:500 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: HCO-18
Immunogen	A synthesized peptide derived from human RUNX2.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	RUNX2
-------------	-------

Synonyms

AML3, CBFA1, OSF2, PEBP2A

Function

Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis (PubMed:[28505335](#), PubMed:[28703881](#), PubMed:[28738062](#), PubMed:[16299379](#)). Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'-PYGPYGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T- cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters. In osteoblasts, supports transcription activation: synergizes with SPEN/MINT to enhance FGFR2-mediated activation of the osteocalcin FGF-responsive element (OCFRE) (By similarity). Inhibits KAT6B-dependent transcriptional activation.


Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q08775}


Tissue Location

Specifically expressed in osteoblasts.

Images



Immunohistochemical analysis of paraffin-embedded human colon, using RUNX2 Antibody.



Immunofluorescent analysis of Saos-2 cells, using RUNX2 Antibody.




Figure 1. Western blot analysis of RUNX2 using anti-RUNX2 antibody (M00442).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human THP-1 whole cell lysates,

Lane 2: human U251 whole cell lysates,

Lane 3: human RT4 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RUNX2 antigen affinity purified monoclonal antibody (Catalog # M00442) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent

detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for RUNX2 at approximately 57 kDa. The expected band size for RUNX2 is at 57 kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.