10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Granzyme B GZMB Monoclonal Antibody

Catalog # ABO14382

Product Information

Application	IHC, IF, ICC, IP
Primary Accession	P10144
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Granzyme B GZMB Monoclonal Antibody . Tested in IHC, ICC/IF, IP applications. This antibody reacts with Human.


Additional Information

Gene ID	3002
Other Names	Granzyme B, 3.4.21.79, C11, CTLA-1, Cathepsin G-like 1, CTSGL1, Cytotoxic T-lymphocyte proteinase 2, Lymphocyte protease, Fragmentin-2, Granzyme-2, Human lymphocyte protein, HLP, SECT, T-cell serine protease 1-3E, GZMB {ECO:0000303|PubMed:32188940, ECO:0000312|HGNC:HGNC:4709}
Calculated MW	27716
Application Details	IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AAHC-7
Immunogen	A synthesized peptide derived from human Granzyme B This enzyme is necessary for target cell lysis in cell-mediated immune responses. It cleaves after Asp. Seems to be linked to an activation cascade of caspases (aspartate-specific cysteine proteases) responsible for apoptosis execution. Cleaves caspase-3, -7, -9 and 10 to give rise to active enzymes mediating apoptosis.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GZMB {ECO:0000303|PubMed:32188940, ECO:0000312|HGNC:HGNC:4709}
Function	Abundant protease in the cytosolic granules of cytotoxic T- cells and NK-cells which activates caspase-independent pyroptosis when delivered into the target cell through the immunological synapse (PubMed:1985927, PubMed:3262682, PubMed:3263427). It cleaves after Asp (PubMed:1985927, PubMed:8258716). Once delivered into the target cell, acts by catalyzing cleavage of gasdermin-E (GSDME), releasing the pore- forming moiety of GSDME, thereby triggering pyroptosis and target cell death (PubMed:31953257, PubMed:32188940). Seems to be linked to an activation cascade of caspases (aspartate-specific cysteine proteases) responsible for apoptosis execution. Cleaves caspase-3, -9 and -10 (CASP3, CASP9 and CASP10, respectively) to give rise to active enzymes mediating apoptosis (PubMed:9852092). Cleaves and activates CASP7 in response to bacterial infection, promoting plasma membrane repair (By similarity).
Cellular Location	Secreted. Cytolytic granule. Note=Delivered into the target cell by perforin (PubMed:20038786).

Images



Immunohistochemical analysis of paraffin-embedded human T cell lymphoma, using Granzyme B Antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.