10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-LMO2 Monoclonal Antibody

Catalog # ABO14388

## Product Information

| | |
|---|---|
| **Application** | WB, IHC, IP, FC |
| **Primary Accession** | P25791 |
| **Host** | Rabbit |
| **Isotype** | Rabbit IgG |
| **Reactivity** | Rat, Human, Mouse |
| **Clonality** | Monoclonal |
| **Format** | Liquid |
| **Description** | Anti-LMO2 Monoclonal Antibody . Tested in WB, IHC, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 4005 |
| **Other Names** | Rhombotin-2, Cysteine-rich protein TTG-2, LIM domain only protein 2, LMO-2, T-cell translocation protein 2, LMO2, RBTN2, RBTNL1, RHOM2, TTG2 |
| **Calculated MW** | 18358 |
| **Application Details** | WB 1:500-1:2000<br>IHC 1:50-1:200<br>IP 1:100<br>FC 1:100 |
| **Source** | Eukaryota |
| **Contents** | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| **Clone Names** | Clone: ACOA-12 |
| **Immunogen** | A synthesized peptide derived from human LMO2 Acts with TAL1/SCL to regulate red blood cell development. Also acts with LDB1 to maintain erythroid precursors in an immature state. |
| **Purification** | Affinity-chromatography |
| **Storage** | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |

## Protein Information

| | |
|---|---|
| **Name** | LMO2 |

| | |
|---|---|
| **Synonyms** | RBTN2, RBTNL1, RHOM2, TTG2 |
| **Function** | Acts with TAL1/SCL to regulate red blood cell development. Also acts with LDB1 to maintain erythroid precursors in an immature state. |
| **Cellular Location** | Nucleus. |

## Images




Figure 1. Western blot analysis of LMO2 using anti-LMO2 antibody (M03502).
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.
Lane 1: human Raji whole cell lysates,
Lane 2: rat C6 whole cell lysates,
Lane 3: mouse Neuro-2a whole cell lysates.
After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-LMO2 antigen affinity purified monoclonal antibody (Catalog # M03502) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for LMO2 at approximately 19 kDa. The expected band size for LMO2 is at 19 kDa.



Figure 2. IHC analysis of LMO2 using anti-LMO2 antibody (M03502).
LMO2 was detected in a paraffin-embedded section of lymph node of rat lung tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LMO2 Antibody (M03502) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Figure 3. IHC analysis of LMO2 using anti-LMO2 antibody (M03502).
LMO2 was detected in a paraffin-embedded section of lymph node of rat lung tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LMO2 Antibody (M03502) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30



minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Figure 4. IHC analysis of LMO2 using anti-LMO2 antibody (M03502).
LMO2 was detected in a paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LMO2 Antibody (M03502) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



Figure 5. IHC analysis of LMO2 using anti-LMO2 antibody (M03502).
LMO2 was detected in a paraffin-embedded section of mouse lung tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-LMO2 Antibody (M03502) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.