

Anti-WASL Monoclonal Antibody

Catalog # ABO14389

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	O00401
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-WASL Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information

Gene ID	8976
Other Names	Actin nucleation-promoting factor WASL, Neural Wiskott-Aldrich syndrome protein, N-WASP, WASL
Calculated MW	54827
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: ACOB-23 A synthesized peptide derived from human WASL Regulates actin polymerization by stimulating the actin-nucleating activity of the Arp2/3 complex. Binds to HSF1/HSTF1 and forms a complex on heat shock promoter elements (HSE) that negatively regulates HSP90 expression.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	WASL
-------------	------


Function

Regulates actin polymerization by stimulating the actin- nucleating activity of the Arp2/3 complex (PubMed:[16767080](#), PubMed:[19366662](#), PubMed:[19487689](#), PubMed:[22847007](#), PubMed:[22921828](#), PubMed:[9422512](#)). Involved in various processes, such as mitosis and cytokinesis, via its role in the regulation of actin polymerization (PubMed:[19366662](#), PubMed:[19487689](#), PubMed:[22847007](#), PubMed:[22921828](#), PubMed:[9422512](#)). Together with CDC42, involved in the extension and maintenance of the formation of thin, actin-rich surface projections called filopodia (PubMed:[9422512](#)). In addition to its role in the cytoplasm, also plays a role in the nucleus by regulating gene transcription, probably by promoting nuclear actin polymerization (PubMed:[16767080](#)). Binds to HSF1/HSTF1 and forms a complex on heat shock promoter elements (HSE) that negatively regulates HSP90 expression (By similarity). Plays a role in dendrite spine morphogenesis (By similarity). Decreasing levels of DNMBP (using antisense RNA) alters apical junction morphology in cultured enterocytes, junctions curve instead of being nearly linear (PubMed:[19767742](#)).

Cellular Location

Cytoplasm, cytoskeleton. Nucleus Cytoplasm {ECO:0000250|UniProtKB:Q91YD9}. Note=Preferentially localized in the cytoplasm when phosphorylated and in the nucleus when unphosphorylated (By similarity). Exported from the nucleus by an nuclear export signal (NES)-dependent mechanism to the cytoplasm (By similarity). {ECO:0000250|UniProtKB:Q91YD9}

Images



Western blot analysis of WASL expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.