10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-PDK1 Monoclonal Antibody

Catalog # ABO14393

Product Information

Application	WB, IP
Primary Accession	Q15118
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-PDK1 Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information


Gene ID	5163
Other Names	[Pyruvate dehydrogenase (acetyl-transferring)] kinase isozyme 1, mitochondrial, 2.7.11.2, Pyruvate dehydrogenase kinase isoform 1, PDH kinase 1, PDK1, PDHK1
Calculated MW	49244
Application Details	WB 1:500-1:2000 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ACOF-16
Immunogen	A synthesized peptide derived from human PDK1 Inhibits the mitochondrial pyruvate dehydrogenase complex by phosphorylation of the E1 alpha subunit, thus contributing to the regulation of glucose metabolism.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	PDK1
Synonyms	PDHK1

Function	Kinase that plays a key role in regulation of glucose and fatty acid metabolism and homeostasis via phosphorylation of the pyruvate dehydrogenase subunits PDHA1 and PDHA2 (PubMed:7499431, PubMed:18541534, PubMed:22195962, PubMed:26942675, PubMed:17683942). This inhibits pyruvate dehydrogenase activity, and thereby regulates metabolite flux through the tricarboxylic acid cycle, down-regulates aerobic respiration and inhibits the formation of acetyl-coenzyme A from pyruvate (PubMed:18541534, PubMed:22195962, PubMed:26942675). Plays an important role in cellular responses to hypoxia and is important for cell proliferation under hypoxia (PubMed:18541534, PubMed:22195962, PubMed:26942675).
Cellular Location	Mitochondrion matrix
Tissue Location	Expressed predominantly in the heart. Detected at lower levels in liver, skeletal muscle and pancreas


Images



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of PDK1 expression in LNCaP cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.