

Anti-CDT1 Rabbit Monoclonal Antibody

Catalog # ABO14398

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	Q9H211
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-CDT1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	81620
Other Names	DNA replication factor Cdt1, Double parked homolog, DUP, CDT1 {ECO:0000312 EMBL:AAG45181.1}
Calculated MW	60390
Application Details	WB 1:500-1:1000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: BO-3 A synthesized peptide derived from human CDT1.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	CDT1 {ECO:0000312 EMBL:AAG45181.1}
Function	Required for both DNA replication and mitosis (PubMed: 11125146 , PubMed: 14993212 , PubMed: 21856198 , PubMed: 22581055 ,

PubMed:[26842564](#)). DNA replication licensing factor, required for pre-replication complex assembly. Cooperates with CDC6 and the origin recognition complex (ORC) during G1 phase of the cell cycle to promote the loading of the mini-chromosome maintenance (MCM) complex onto DNA to generate pre-replication complexes (pre-RC) (PubMed:[14672932](#)). Required also for mitosis by promoting stable kinetochore-microtubule attachments (PubMed:[22581055](#)). Potential oncogene (By similarity).

Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Note=Transiently localizes to kinetochores during prometaphase and metaphase

Images



Western blot analysis of CDT1 expression in HEK293 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.