

Anti-CEACAM6 Rabbit Monoclonal Antibody

Catalog # ABO14411

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	P40199
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CEACAM6 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4680
Other Names	Carcinoembryonic antigen-related cell adhesion molecule 6, Non-specific crossreacting antigen, Normal cross-reacting antigen, CD66c, CEACAM6, NCA
Calculated MW	37237
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ACBC-3
Immunogen	A synthesized peptide derived from human CEACAM6
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CEACAM6 (HGNC:1818)
Function	Cell surface glycoprotein that plays a role in cell adhesion and tumor progression (PubMed: 10910050 , PubMed: 11590190 , PubMed: 1378450 ,

PubMed:[16204051](#), PubMed:[2022629](#), PubMed:[2803308](#), PubMed:[8776764](#)). Intercellular adhesion occurs in a calcium- and fibronectin-independent manner (PubMed:[16204051](#), PubMed:[2022629](#)). Mediates homophilic and heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM5 and CEACAM8 (PubMed:[11590190](#), PubMed:[16204051](#), PubMed:[2022629](#), PubMed:[2803308](#), PubMed:[8776764](#)). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed:[8776764](#)). Plays a role in neutrophil adhesion to cytokine-activated endothelial cells (PubMed:[1378450](#)). Plays a role in cell migration and cell adhesion to endothelial cells (PubMed:[16204051](#)).


Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor. Apical cell membrane. Cell surface. Note=Localized to the apical glycocalyx surface.


Tissue Location

Expressed in neutrophils (PubMed:1378450). Expressed in columnar epithelial and goblet cells of the colon (PubMed:10436421). Expressed in numerous tumor cell lines (at protein level) (PubMed:16204051).

Images



Western blot analysis of CEACAM6 expression in MCF7 cell lysate.



All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.