

Anti-Factor H CFH Monoclonal Antibody

Catalog # ABO14427

Product Information

Application	WB, IF, ICC
Primary Accession	P08603
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Factor H CFH Monoclonal Antibody . Tested in WB, ICC/IF applications. This antibody reacts with Human.

Additional Information


Gene ID	3075
Other Names	Complement factor H, H factor 1, CFH, HF, HF1, HF2
Calculated MW	139096
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: ACCI-3 A synthesized peptide derived from human Factor H Factor H functions as a cofactor in the inactivation of C3b by factor I and also increases the rate of dissociation of the C3bBb complex (C3 convertase) and the (C3b) NBB complex (C5 convertase) in the alternative complement pathway.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CFH
Synonyms	HF, HF1, HF2

Function	Glycoprotein that plays an essential role in maintaining a well-balanced immune response by modulating complement activation. Acts as a soluble inhibitor of complement, where its binding to self markers such as glycan structures prevents complement activation and amplification on cell surfaces (PubMed: 21285368 , PubMed: 21317894 , PubMed: 25402769). Accelerates the decay of the complement alternative pathway (AP) C3 convertase C3bBb, thus preventing local formation of more C3b, the central player of the complement amplification loop (PubMed: 19503104 , PubMed: 21317894 , PubMed: 26700768). As a cofactor of the serine protease factor I, CFH also regulates proteolytic degradation of already-deposited C3b (PubMed: 18252712 , PubMed: 23332154 , PubMed: 28671664). In addition, mediates several cellular responses through interaction with specific receptors. For example, interacts with CR3/ITGAM receptor and thereby mediates the adhesion of human neutrophils to different pathogens. In turn, these pathogens are phagocytosed and destroyed (PubMed: 20008295 , PubMed: 9558116).
Cellular Location	Secreted.
Tissue Location	Expressed in the retinal pigment epithelium (at protein level) (PubMed: 25136834). CFH is one of the most abundant complement components in blood where the liver is the major source of CFH protein in vivo. In addition, CFH is secreted by additional cell types including monocytes, fibroblasts, or endothelial cells (PubMed: 2139673 , PubMed: 25136834 , PubMed: 2968404 , PubMed: 6444659).

Images



Western blot analysis of Factor H expression in human plasma lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.