

Anti-CXCR3 Monoclonal Antibody

Catalog # ABO14451

Product Information

Application	WB, IF, ICC, IP
Primary Accession	P49682
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-CXCR3 Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human.


Additional Information

Gene ID	2833
Other Names	C-X-C chemokine receptor type 3, CXC-R3, CXCR-3, CKR-L2, G protein-coupled receptor 9, Interferon-inducible protein 10 receptor, IP-10 receptor, CD183, CXCR3, GPR9
Calculated MW	40660
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: ACFD-3 A synthesized peptide derived from human CXCR3 Receptor for CXCL9, CXCL10 and CXCL11 and mediates the proliferation of human mesangial cells (HMC). Isoform 2 is a receptor for CXCL4 and also mediates the inhibitory activities of CXCL9, CXCL10 and CXCL11 on the growth of human microvascular endothelial cells (HMVEC). May play a role in angiogenesis.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CXCR3
Synonyms	GPR9
Function	[Isoform 1]: Receptor for the C-X-C chemokines CXCL9, CXCL10 and CXCL11, and mediates the proliferation, survival and angiogenic activity of human mesangial cells (HMC) through a heterotrimeric G- protein signaling pathway (PubMed: 12782716). Probably promotes cell chemotaxis response. Upon activation by PF4, induces activated T- lymphocyte migration mediated via downstream Ras/extracellular signal- regulated kinase (ERK) signaling. [Isoform 3]: Mediates the activity of CXCL11.
Cellular Location	[Isoform 1]: Cell membrane; Multi-pass membrane protein
Tissue Location	Isoform 1 and isoform 2 are mainly expressed in heart, kidney, liver and skeletal muscle. Isoform 1 is also expressed in placenta. Isoform 2 is expressed in endothelial cells. Expressed in T-cells (at protein level).

Images



Western blot analysis of CXCR3 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.