

Anti-DNA Polymerase beta Monoclonal Antibody

Catalog # ABO14465

Product Information

Application	WB, IHC, IF, ICC, IP
Primary Accession	P06746
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-DNA Polymerase beta Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	5423
Other Names	DNA polymerase beta, 2.7.7.7, 5'-deoxyribose-phosphate lyase, 5'-dRP lyase, 4.2.99.-, AP lyase, 4.2.99.18, POLB
Calculated MW	38178
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: ACGH-16 A synthesized peptide derived from human DNA Polymerase beta Repair polymerase. Conducts "gap-filling" DNA synthesis in a stepwise distributive fashion rather than in a processive fashion as for other DNA polymerases. Has a 5'-deoxyribose-5-phosphate lyase (dRP lyase) activity.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	POLB
-------------	------


Function

Repair polymerase that plays a key role in base-excision repair (PubMed:[10556592](#), PubMed:[9207062](#), PubMed:[9572863](#)). During this process, the damaged base is excised by specific DNA glycosylases, the DNA backbone is nicked at the abasic site by an apurinic/apyrimidic (AP) endonuclease, and POLB removes 5'-deoxyribose-phosphate from the preincised AP site acting as a 5'-deoxyribose-phosphate lyase (5'-dRP lyase); through its DNA polymerase activity, it adds one nucleotide to the 3' end of the arising single-nucleotide gap (PubMed:[10556592](#), PubMed:[17526740](#), PubMed:[8841119](#), PubMed:[9556598](#), PubMed:[9572863](#), PubMed:[9614142](#)). Conducts 'gap-filling' DNA synthesis in a stepwise distributive fashion rather than in a processive fashion as for other DNA polymerases. It is also able to cleave sugar-phosphate bonds 3' to an intact AP site, acting as an AP lyase (PubMed:[9614142](#)).


Cellular Location

Nucleus. Cytoplasm. Note=Cytoplasmic in normal conditions. Translocates to the nucleus following DNA damage


Images



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.



Western blot analysis of DNA Polymerase beta expression in A431 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.