

Anti-TRPM7 Monoclonal Antibody

Catalog # ABO14560

Product Information

Application	WB, IF, ICC, FC
Primary Accession	Q96QT4
Host	Rabbit
Isotype	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-TRPM7 Monoclonal Antibody . Tested in WB, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	54822
Other Names	Transient receptor potential cation channel subfamily M member 7, 2.7.11.1, Channel-kinase 1, Long transient receptor potential channel 7, LTrpC-7, LTrpC7, TRPM7, CHAK1, LTRPC7
Calculated MW	212697
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 FC 1:100
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: ADGD-20 A synthesized peptide derived from human TRPM7
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	TRPM7
Synonyms	CHAK1, LTRPC7 {ECO:0000303 PubMed:113855}


Function

Bifunctional protein that combines an ion channel with an intrinsic kinase domain, enabling it to modulate cellular functions either by conducting ions through the pore or by phosphorylating downstream proteins via its kinase domain. The channel is highly permeable to divalent cations, specifically calcium (Ca^{2+}), magnesium (Mg^{2+}) and zinc (Zn^{2+}) and mediates their influx (PubMed:[11385574](#), PubMed:[12887921](#), PubMed:[15485879](#), PubMed:[24316671](#), PubMed:[35561741](#), PubMed:[36027648](#)). Controls a wide range of biological processes such as Ca^{2+} , Mg^{2+} and Zn^{2+} homeostasis, vesicular Zn^{2+} release channel and intracellular Ca^{2+} signaling, embryonic development, immune responses, cell motility, proliferation and differentiation (By similarity). The C-terminal alpha-kinase domain autophosphorylates cytoplasmic residues of TRPM7 (PubMed:[18365021](#)). In vivo, TRPM7 phosphorylates SMAD2, suggesting that TRPM7 kinase may play a role in activating SMAD signaling pathways. In vitro, TRPM7 kinase phosphorylates ANXA1 (annexin A1), myosin II isoforms and a variety of proteins with diverse cellular functions (PubMed:[15485879](#), PubMed:[18394644](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q923J1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Note=Localized largely in intracellular Zn^{2+} -storage vesicles. {ECO:0000250|UniProtKB:Q923J1}

Images



Western blot analysis of TRPM7 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.