

Anti-ANP NPPA Monoclonal Antibody

Catalog # ABO14570

Product Information

Application	WB, IP
Primary Accession	P01160
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-ANP NPPA Monoclonal Antibody . Tested in WB, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	4878
Other Names	Natriuretic peptides A, Atrial natriuretic factor prohormone, proANF, Atrial natriuretic peptide prohormone, preproANP, proANP, Atriopeptigen, Cardiodilatin, CDD, preproCDD-ANF, Long-acting natriuretic peptide, LANP, Long-acting natriuretic hormone, LANH, Pro atrial natriuretic factor 1-30, proANF 1-30, Pro atrial natriuretic peptide 1-30, proANP 1-30, Vessel dilator, VSDL, Pro atrial natriuretic factor 31-67, proANF 31-67, Pro atrial natriuretic peptide 31-67, proANP 31-67, Kaliuretic peptide, KP, Pro atrial natriuretic factor 79-98, proANF 79-98, Pro atrial natriuretic peptide 79-98, proANP 79-98, Urodilatin, URO, CDD 95-126, CDD-ANP (95-126), Pro atrial natriuretic peptide 95-126, proANP 95-126, Auriculin-C, Atrial natriuretic factor 1-33, NPPA, ANP, PND
Calculated MW	16396
Application Details	WB 1:500-1:2000 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: ADHD-14
Immunogen	A synthesized peptide derived from human ANP Hormone playing a key role in cardiovascular homeostasis through regulation of natriuresis, diuresis, and vasodilation. Also plays a role in female pregnancy by promoting trophoblast invasion and spiral artery remodeling in uterus.
Purification	Affinity-chromatography

Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
----------------	--

Protein Information

Name	NPPA
-------------	------

Synonyms	ANP, PND
-----------------	----------


Function	[Atrial natriuretic peptide]: Hormone that plays a key role in mediating cardio-renal homeostasis, and is involved in vascular remodeling and regulating energy metabolism (PubMed:15741263, PubMed:16875975, PubMed:18835931, PubMed:21672517, PubMed:22307324, PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Acts by specifically binding and stimulating NPR1 to produce cGMP, which in turn activates effector proteins, such as PRKG1, that drive various biological responses (PubMed:1660465, PubMed:1672777, PubMed:21098034, PubMed:2162527, PubMed:22307324, PubMed:2825692, PubMed:7720651, PubMed:8384600, PubMed:9893117). Regulates vasodilation, natriuresis, diuresis and aldosterone synthesis and is therefore essential for regulating blood pressure, controlling the extracellular fluid volume and maintaining the fluid-electrolyte balance (PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Also involved in inhibiting cardiac remodeling and cardiac hypertrophy by inducing cardiomyocyte apoptosis and attenuating the growth of cardiomyocytes and fibroblasts (PubMed:16875975). Plays a role in female pregnancy by promoting trophoblast invasion and spiral artery remodeling in uterus, and thus prevents pregnancy-induced hypertension (By similarity). In adipose tissue, acts in various cGMP- and PKG- dependent pathways to regulate lipid metabolism and energy homeostasis (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). This includes up-regulating lipid metabolism and mitochondrial oxygen utilization by activating the AMP-activated protein kinase (AMPK), and increasing energy expenditure by acting via MAPK11 to promote the UCP1- dependent thermogenesis of brown adipose tissue (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). Binds the clearance receptor NPR3 which removes the hormone from circulation (PubMed:1672777).
-----------------	---

Cellular Location	[Long-acting natriuretic peptide]: Secreted. Note=Detected in blood. [Kaliuretic peptide]: Secreted. Note=Detected in blood [Atrial natriuretic peptide]: Secreted. Perikaryon. Cell projection. Note=Detected in blood (PubMed:15741263, PubMed:18835931, PubMed:2532366, PubMed:7955907, PubMed:7984506, PubMed:8351194, PubMed:8653797, PubMed:8779891). Detected in urine in one study (PubMed:8351194). However, in another study, was not detected in urine (PubMed:7984506). Detected in cytoplasmic bodies and neuronal processes of pyramidal neurons (layers II-VI) (PubMed:30534047) Increased secretion in response to the vasopressin AVP (By similarity) Likely to be secreted in response to an increase in atrial pressure or atrial stretch (PubMed:2532366). In kidney cells, secretion increases in response to activated guanylyl cyclases and increased intracellular cAMP levels (PubMed:9893117). Plasma levels increase 15 minutes after a high-salt meal, and decrease back to normal plasma levels 1 hr later (PubMed:8779891). {ECO:0000250 UniProtKB:P01161, ECO:0000269 PubMed:15741263, ECO:0000269 PubMed:18835931, ECO:0000269 PubMed:2532366, ECO:0000269 PubMed:30534047, ECO:0000269 PubMed:7955907, ECO:0000269 PubMed:7984506, ECO:0000269 PubMed:8351194, ECO:0000269 PubMed:8653797, ECO:0000269 PubMed:8779891, ECO:0000269 PubMed:9893117}
--------------------------	--

Tissue Location

[Urodilatin]: Detected in the kidney distal tubular cells (at protein level) (PubMed:8384600, PubMed:9794555). Present in urine (at protein level) (PubMed:2972874, PubMed:8351194, PubMed:8779891, PubMed:9794555).

Images



Western blot analysis of ANP expression in PC3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.