10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-NuMA NUMA1 Monoclonal Antibody

Catalog # ABO14613

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	Q14980
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-NuMA NUMA1 Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	4926
Other Names	Nuclear mitotic apparatus protein 1 {ECO:0000312|HGNC:HGNC:8059}, Nuclear matrix protein-22, NMP-22, Nuclear mitotic apparatus protein, NuMA protein, SP-H antigen, NUMA1 (HGNC:8059)
Calculated MW	238260
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AEBF-14
Immunogen	A synthesized peptide derived from human NuMA
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	NUMA1 (HGNC:8059)
Function	Microtubule (MT)-binding protein that plays a role in the formation and maintenance of the spindle poles and the alignement and the segregation of


chromosomes during mitotic cell division (PubMed:17172455, PubMed:19255246, PubMed:24996901, PubMed:26195665, PubMed:27462074, PubMed:7769006). Functions to tether the minus ends of MTs at the spindle poles, which is critical for the establishment and maintenance of the spindle poles (PubMed:11956313, PubMed:12445386). Plays a role in the establishment of the mitotic spindle orientation during metaphase and elongation during anaphase in a dynein-dynactin- dependent manner (PubMed:23870127, PubMed:24109598, PubMed:24996901, PubMed:26765568). In metaphase, part of a ternary complex composed of GPSM2 and G(i) alpha proteins, that regulates the recruitment and anchorage of the dynein-dynactin complex in the mitotic cell cortex regions situated above the two spindle poles, and hence regulates the correct oritentation of the mitotic spindle (PubMed:22327364, PubMed:23027904, PubMed:23921553). During anaphase, mediates the recruitment and accumulation of the dynein-dynactin complex at the cell membrane of the polar cortical region through direct association with phosphatidylinositol 4,5-bisphosphate (PI(4,5)P2), and hence participates in the regulation of the spindle elongation and chromosome segregation (PubMed:22327364, PubMed:23921553, PubMed:24371089, PubMed:24996901). Also binds to other polyanionic phosphoinositides, such as phosphatidylinositol 3-phosphate (PIP), lysophosphatidic acid (LPA) and phosphatidylinositol triphosphate (PIP3), in vitro (PubMed:24371089, PubMed:24996901). Also required for proper orientation of the mitotic spindle during asymmetric cell divisions (PubMed:21816348). Plays a role in mitotic MT aster assembly (PubMed:11163243, PubMed:11229403, PubMed:12445386). Involved in anastral spindle assembly (PubMed:25657325). Positively regulates TNKS protein localization to spindle poles in mitosis (PubMed:16076287). Highly abundant component of the nuclear matrix where it may serve a non-mitotic structural role, occupies the majority of the nuclear volume (PubMed:10075938). Required for epidermal differentiation and hair follicle morphogenesis (By similarity).

Cellular Location

Nucleus. Nucleus, nucleoplasm. Nucleus matrix. Chromosome. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cell cortex. Cell membrane; Lipid-anchor; Cytoplasmic side. Lateral cell membrane {ECO:0000250|UniProtKB:E9Q7G0}. Note=Mitotic cell cycle- dependent shuttling protein that relocalizes from the interphase nucleus to the spindle poles and cell cortex (PubMed:10811826, PubMed:1541636). The localization to the spindle poles is regulated by AAAS (PubMed:26246606). In interphase, resides in the nuclear matrix (PubMed:1541630, PubMed:1541636, PubMed:23921553). In prophase, restricted to the interchromatin or condensed chromosome space (PubMed:10811826). In prometaphase, after nuclear envelope disassembly, forms aggregates both in the spindle midzone and at duplicated centrosomes and astral microtubules (MTs) of the bipolar spindle apparatus (PubMed:10811826). Translocates from the spindle midzone towards the spindle poles along spindle fibers in a MT- and dynein-dynactin-dependent manner until the anaphase onset (PubMed:10811826, PubMed:1541636). In metaphase, recruited to the polar cortical region in a GPSM2- and GNAI1-dependent manner (PubMed:23870127, PubMed:24109598, PubMed:24996901). Excluded from the metaphase equatorial cortical region in a RanGTP-dependent manner (PubMed:22327364, PubMed:23870127). Phosphorylation on Thr-2055 by CDK1 results in its localization at spindle poles in metaphase, but not at the cell cortex (PubMed:23921553). In anaphase, recruited and anchored at the cell membrane of the polar cortical region in a EPB41-, EPB41L2-, phosphatidylinositol-dependent and GPSM2- and G(i) alpha proteins-independent manner (PubMed:23870127, PubMed:24109598, PubMed:24371089, PubMed:24996901). Excluded from the anaphase equatorial region of the cell cortex in a RACGAP1- and KIF23-dependent and

RanGTP-independent manner (PubMed:24996901). Associated with astral MTs emanating from the spindle poles during anaphase (PubMed:12445386, PubMed:24996901). Nonphosphorylated Thr-2055 localizes at the cell cortex, weakly during metaphase and more prominently during anaphase in a phosphatase PPP2CA-dependent manner (PubMed:23921553). As mitosis progresses it reassociates with telophase chromosomes very early during nuclear reformation, before substantial accumulation of lamins on chromosomal surfaces is evident (PubMed:1541636). Localizes to the tips of cortical MTs in prometaphase (PubMed:26765568). Localizes along MTs and specifically to both MT plus and minus ends (PubMed:26765568). Also accumulates at MT tips near the cell periphery (PubMed:26765568) Colocalizes with GPSM2 at mitotic spindle poles during mitosis (PubMed:11781568, PubMed:21816348). Colocalizes with SPAG5 at mitotic spindle at prometaphase and at mitotic spindle poles at metaphase and anaphase (PubMed:27462074). Colocalizes with ABRO1 at mitotic spindle poles (PubMed:26195665). Colocalized with TNKS from prophase through to anaphase in mitosis (PubMed:16076287). Colocalizes with tubulin alpha (PubMed:12445386). CCSAP is essential for its centrosomal localization (PubMed:26562023). In horizontally retinal progenitor dividing cells, localized to the lateral cortical region (By similarity) {ECO:0000250|UniProtKB:E9Q7G0, ECO:0000269|PubMed:10811826, ECO:0000269|PubMed:11781568, ECO:0000269|PubMed:12445386, ECO:0000269|PubMed:1541630, ECO:0000269|PubMed:1541636, ECO:0000269|PubMed:16076287, ECO:0000269|PubMed:21816348, ECO:0000269|PubMed:22327364, ECO:0000269|PubMed:23870127, ECO:0000269|PubMed:23921553, ECO:0000269|PubMed:24109598, ECO:0000269|PubMed:24371089, ECO:0000269|PubMed:24996901, ECO:0000269|PubMed:26195665, ECO:0000269|PubMed:26246606, ECO:0000269|PubMed:26562023, ECO:0000269|PubMed:26765568, ECO:0000269|PubMed:27462074} [Isoform 4]: Cytoplasm, cytosol. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Note=During interphase, mainly clustered at the centrosomal region in the cytosol After entry into mitosis, detected at mitotic spindle poles

Images



Western blot analysis of Mad2L1 expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.