

Anti-Geminin Monoclonal Antibody

Catalog # ABO14657

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	O75496
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Geminin Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	51053
Other Names	Geminin, GMNN
Calculated MW	23565
Application Details	WB 1:500-1:1000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AEGO-7
Immunogen	A synthesized peptide derived from human Geminin Inhibits DNA replication by preventing the incorporation of MCM complex into prereplication complex (pre-RC). It is degraded during the mitotic phase of the cell cycle. Its destruction at the metaphase-anaphase transition permits replication in the succeeding cell cycle.
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	GMNN
-------------	------


Function

Inhibits DNA replication by preventing the incorporation of MCM complex into pre-replication complex (pre-RC) (PubMed:[14993212](#), PubMed:[20129055](#), PubMed:[24064211](#), PubMed:[9635433](#)). It is degraded during the mitotic phase of the cell cycle (PubMed:[14993212](#), PubMed:[24064211](#), PubMed:[9635433](#)). Its destruction at the metaphase- anaphase transition permits replication in the succeeding cell cycle (PubMed:[14993212](#), PubMed:[24064211](#), PubMed:[9635433](#)). Inhibits histone acetyltransferase activity of KAT7/HBO1 in a CDT1-dependent manner, inhibiting histone H4 acetylation and DNA replication licensing (PubMed:[20129055](#)). Inhibits the transcriptional activity of a subset of Hox proteins, enrolling them in cell proliferative control (PubMed:[22615398](#)).

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic but can be relocalized to the nucleus.

Images



Western blot analysis of Geminin expression in HeLa cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.