

Anti-KDM4A Monoclonal Antibody

Catalog # ABO14668

Product Information

Application	WB, IHC, IP
Primary Accession	O75164
Host	Rabbit
Isotype	Rabbit IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-KDM4A Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	9682
Other Names	Lysine-specific demethylase 4A, 1.14.11.66, 1.14.11.69, JmjC domain-containing histone demethylation protein 3A, Jumonji domain-containing protein 2A, [histone H3]-trimethyl-L-lysine(36) demethylase 4A, [histone H3]-trimethyl-L-lysine(9) demethylase 4A, KDM4A, JHDM3A, JMJD2, JMJD2A, KIAA0677
Calculated MW	120662
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 IP 1:60
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: AEHB-11
Immunogen	A synthesized peptide derived from human KDM4A
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	KDM4A
-------------	-------

Synonyms

JHDM3A, JMJD2, JMJD2A, KIAA0677

Function

Histone demethylase that specifically demethylates 'Lys-9' and 'Lys-36' residues of histone H3, thereby playing a central role in histone code (PubMed:[16603238](#), PubMed:[26741168](#), PubMed:[21768309](#)). Does not demethylate histone H3 'Lys-4', H3 'Lys-27' nor H4 'Lys-20' (PubMed:[16603238](#), PubMed:[26741168](#), PubMed:[21768309](#)). Demethylates trimethylated H3 'Lys-9' and H3 'Lys-36' residue, while it has no activity on mono- and dimethylated residues (PubMed:[16603238](#), PubMed:[26741168](#), PubMed:[21768309](#)). Demethylation of Lys residue generates formaldehyde and succinate (PubMed:[16603238](#)). Also able to demethylate histone H1-4 methylated at 'Lys-26' (H1.4K26me1, H1.4K26me2 and H1.4K26me3) (PubMed:[19144645](#), PubMed:[30156264](#)). Participates in transcriptional repression of ASCL2 and E2F-responsive promoters via the recruitment of histone deacetylases and NCOR1, respectively (PubMed:[16024779](#)).


Cellular Location

Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00537, ECO:0000269 | PubMed:15927959, ECO:0000269 | PubMed:16024779}


Tissue Location

Ubiquitous..

Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of KDM4A / JHDM3A / JMJD2A expression in (1) HEK293 cell lysate; (2) NIH3T3 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.