

Anti-ADK Antibody Picoband™ (monoclonal, 7F4)

Catalog # ABO14863

Product Information

Application	WB, IHC, FC
Primary Accession	P55263
Host	Mouse
Isotype	Mouse IgG1
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized
Description	Anti-ADK Antibody Picoband™ (monoclonal, 7F4) . Tested in Flow Cytometry, IHC, WB applications. This antibody reacts with Human.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500 µg/ml.

Additional Information

Gene ID	132
Other Names	Adenosine kinase, AK, 2.7.1.20, Adenosine 5'-phosphotransferase, ADK (HGNC:257)
Calculated MW	40545
Application Details	Western blot, 0.1-0.5 µg/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1 µg/ml Flow Cytometry, 1-3 µg/1x10 ⁶ cells
Subcellular Localization	Nucleus. Cytoplasm.
Tissue Specificity	Widely expressed. Highest level in placenta, liver, muscle and kidney.
Source	Eukaryota
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Clone Names Immunogen	Clone: 7F4 E. coli-derived human ADK recombinant protein (Position: K165-T351). Human ADK shares 88.8% and 88.2% amino acid (aa) sequence identity with mouse and rat ADK, respectively.
Cross Reactivity	No cross-reactivity with other proteins.
Storage	Store at -20° C for one year from date of receipt. After reconstitution, at 4° C for one month. It can also be aliquotted and stored frozen at -20° C for six

months. Avoid repeated freeze-thaw cycles.

Protein Information

Name	ADK {ECO:0000303 PubMed:19635462, ECO:0000312 HGNC:HGNC:257}
Function	Adenosine kinase that mediates the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner: catalyzes phosphorylation of both unmodified and modified adenosines (PubMed: 21963049 , PubMed: 40840445 , PubMed: 6246102 , PubMed: 8577746 , PubMed: 9070863). Plays a key role in the detoxification of modified adenosines containing N(6)-methylated adenine (m6A) post- transcriptional modification (PubMed: 40840445). Modified nucleosides are derived from the degradation of RNAs (mRNAs, rRNAs and tRNAs) and possess intrinsic cytotoxicity and must be cleared to prevent metabolic dysfunction (PubMed: 40840445). Catalyzes the phosphorylation of the free cytosolic methylated adenosine nucleotides N(6)-methyladenosine (m6A), N(6),N(6)-dimethyladenosine (m6,6A) and N(6)- isopentenyladenosine (i6A) into adenosine monophosphate (AMP) intermediates that are further detoxified by MAPDA/ADAL (PubMed: 40840445).
Cellular Location	Cytoplasm, cytosol. [Isoform 2]: Cytoplasm
Tissue Location	Widely expressed. Highest level in placenta, liver, muscle and kidney.

Background

This gene is an enzyme which catalyzes the transfer of the gamma-phosphate from ATP to adenosine, thereby serving as a regulator of concentrations of both extracellular adenosine and intracellular adenine nucleotides. Adenosine has widespread effects on the cardiovascular, nervous, respiratory, and immune systems and inhibitors of the enzyme could play an important pharmacological role in increasing intravascular adenosine concentrations and acting as anti-inflammatory agents. Multiple transcript variants encoding different isoforms have been found for this gene.

Images




Figure 1. Western blot analysis of ADK using anti-ADK antibody (M02193). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human PC-3 whole cell lysates, Lane 2: human U-87MG whole cell lysates, Lane 3: human SW620 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-ADK antigen affinity purified monoclonal antibody (Catalog # M02193) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200

system. A specific band was detected for ADK at approximately 45KD. The expected band size for ADK is at 45KD.




Figure 2. IHC analysis of ADK using anti-ADK antibody (M02193).

ADK was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu\text{g/ml}$ mouse anti-ADK Antibody (M02193) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of ADK using anti-ADK antibody (M02193).

ADK was detected in paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu\text{g/ml}$ mouse anti-ADK Antibody (M02193) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of ADK using anti-ADK antibody (M02193).

ADK was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu\text{g/ml}$ mouse anti-ADK Antibody (M02193) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 5. Flow Cytometry analysis of PC-3 cells using anti-ADK antibody (M02193).

Overlay histogram showing PC-3 cells stained with M02193 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-ADK Antibody (M02193, 1 $\mu\text{g}/1 \times 10^6$ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 $\mu\text{g}/1 \times 10^6$ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 $\mu\text{g}/1 \times 10^6$) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.