

Anti-GSDMD Antibody Picoband™ (monoclonal, 3D5)

Catalog # ABO15067

Product Information

Application	WB, IHC
Primary Accession	P57764
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized
Description	Anti-GSDMD Antibody Picoband™ (monoclonal, 3D5) . Tested in IHC, WB applications. This antibody reacts with Human.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Additional Information

Gene ID	79792
Other Names	Gasdermin-D, Gasdermin domain-containing protein 1, Gasdermin-D, N-terminal, GSDMD-NT, Gasdermin-D, 13 kDa, 13 kDa GSDMD, Gasdermin-D, p40, GSDMD {ECO:0000303 PubMed:26375003, ECO:0000312 HGNC:HGNC:25697}
Calculated MW	52801
Application Details	Western blot, 0.25-0.5 µg/ml, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 µg/ml, Human
Source	Eukaryota
Contents	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.
Clone Names	Clone: 3D5
Immunogen	E.coli-derived human GSDMD recombinant protein (Position: M1-H484).
Purification	Immunogen affinity purified.
Storage	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.

Protein Information

Name	GSDMD {ECO:0000303 PubMed:26375003,
-------------	-------------------------------------

Function

[Gasdermin-D]: Precursor of a pore-forming protein that plays a key role in host defense against pathogen infection and danger signals (PubMed:[26375003](#), PubMed:[26375259](#), PubMed:[27281216](#)). This form constitutes the precursor of the pore-forming protein: upon cleavage, the released N-terminal moiety (Gasdermin-D, N-terminal) binds to membranes and forms pores, triggering pyroptosis (PubMed:[26375003](#), PubMed:[26375259](#), PubMed:[27281216](#), PubMed:[40118151](#)).

Cellular Location

[Gasdermin-D]: Cytoplasm, cytosol. Inflammasome {ECO:0000250 | UniProtKB:Q9D8T2}. Note=In response to a canonical inflammasome stimulus, such as nigericin, recruited to NLRP3 inflammasome with similar kinetics to that of uncleaved CASP1 precursor. {ECO:0000250 | UniProtKB:Q9D8T2} [Gasdermin-D, N-terminal]: Cytoplasm, cytosol. Note=(Microbial infection) Upon infection by *M.tuberculosis*, localization to cell membrane is prevented by *M.tuberculosis* phosphatase PtpB that catalyzes dephosphorylation of phosphatidylinositol (4,5)-bisphosphate and phosphatidylinositol 4- phosphate, thereby inhibiting the membrane targeting of Gasdermin-D, N- terminal and subsequent cytokine release and pyroptosis [Gasdermin-D, C-terminal]: Cytoplasm, cytosol {ECO:0000250 | UniProtKB:Q9D8T2}

Tissue Location

Expressed in the suprabasal cells of esophagus, as well as in the isthmus/neck, pit, and gland of the stomach, suggesting preferential expression in differentiating cells

Background

Gasdermin D is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. Gasdermin D has been suggested to act as a tumor suppressor. Alternatively spliced transcript variants have been described.


Images

Figure 1. Western blot analysis of GSDMD using anti-GSDMD antibody (M02842-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human THP-1 whole cell lysates, Lane 2: human SW620 whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human U87 whole cell lysates, Lane 5: human PC-3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-GSDMD antigen affinity purified monoclonal antibody (Catalog # M02842-1) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001)

with Tanon 5200 system. A specific band was detected for GSDMD at approximately 53 kDa. The expected band size for GSDMD is at 53 kDa.




Figure 2. IHC analysis of GSDMD using anti-GSDMD antibody (M02842-1). GSDMD was detected in a paraffin-embedded section of human esophagus tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-GSDMD Antibody (M02842-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of GSDMD using anti-GSDMD antibody (M02842-1). GSDMD was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-GSDMD Antibody (M02842-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of GSDMD using anti-GSDMD antibody (M02842-1). GSDMD was detected in a paraffin-embedded section of human lymph nodes tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 μ g/ml mouse anti-GSDMD Antibody (M02842-1) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.