

Anti-GNG2 Antibody Picoband™ (monoclonal, 7C13)

Catalog # ABO15090

Product Information

Application	WB, IF, ICC
Primary Accession	P59768
Host	Mouse
Isotype	Mouse IgG2b
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Lyophilized
Description	Anti-GNG2 Antibody Picoband™ (monoclonal, 7C13) . Tested in IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.

Additional Information

Gene ID	54331
Other Names	Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2, G gamma-I, GNG2
Calculated MW	7850
Application Details	Western blot, 0.25-0.5 µg/ml, Mouse, Rat Immunocytochemistry/Immunofluorescence, 5 µg/ml, Human
Source	Eukaryota
Contents	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.
Clone Names	Clone: 7C13
Immunogen	E.coli-derived human GNG2 recombinant protein (Position: A2-D48).
Purification	Immunogen affinity purified.
Storage	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.

Protein Information

Name	GNG2
Function	Guanine nucleotide-binding proteins (G proteins) are involved as a

modulator or transducer in various transmembrane signaling systems (PubMed:[29925951](#), PubMed:[33762731](#), PubMed:[34239069](#), PubMed:[35610220](#), PubMed:[35714614](#), PubMed:[35835867](#), PubMed:[36087581](#), PubMed:[36989299](#), PubMed:[37327704](#), PubMed:[37935376](#), PubMed:[37935377](#), PubMed:[37963465](#), PubMed:[38168118](#), PubMed:[38552625](#), PubMed:[39103320](#)). The beta and gamma chains are required for the GTPase activity, for replacement of GDP by GTP, and for G protein-effector interaction (PubMed:[29925951](#), PubMed:[33762731](#), PubMed:[34239069](#), PubMed:[35610220](#), PubMed:[35714614](#), PubMed:[35835867](#), PubMed:[36087581](#), PubMed:[36989299](#), PubMed:[37327704](#), PubMed:[37935376](#), PubMed:[37935377](#), PubMed:[37963465](#), PubMed:[38168118](#), PubMed:[38552625](#), PubMed:[39103320](#)).

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side

Tissue Location

Expressed in fetal tissues, including testis, adrenal gland, brain, white blood cells and brain

Background

Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2 is a protein that in humans is encoded by the GNG2 gene. This gene encodes one of the gamma subunits of a guanine nucleotide-binding protein. Such proteins are involved in signaling mechanisms across membranes. Various subunits forms heterodimers which then interact with the different signal molecules.

Images




Figure 1. Western blot analysis of GNG2 using anti-GNG2 antibody (M06975-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.


Lane 1: rat brain tissue lysates,

Lane 2: mouse brain tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-GNG2 antigen affinity purified monoclonal antibody (Catalog # M06975-1) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for GNG2 at approximately 12 kDa. The expected band size for GNG2 is at 8 kDa.

Figure 2. IF analysis of GNG2 using anti-GNG2 antibody (M06975-1).

GNG2 was detected in an immunocytochemical section of Caco-2 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum.



And then incubated with 5 µg/mL mouse anti-GNG2 Antibody (M06975-1) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.