10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8)

Catalog # ABO15095

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	P13647
Host	Mouse
Isotype	Mouse IgG2a
Reactivity	Human
Clonality	Monoclonal
Format	Lyophilized
Description	Anti-Cytokeratin 5 Antibody Picoband™ (monoclonal, 9H2F8) . Tested in FCM, IF, IHC, ICC, WB applications. This antibody reacts with Human.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 □g/ml.

Additional Information

Gene ID	3852
Other Names	Keratin, type II cytoskeletal 5, 58 kDa cytokeratin, Cytokeratin-5, CK-5, Keratin-5, K5, Type-II keratin Kb5, KRT5
Calculated MW	62378
Application Details	Western blot, 0.25-0.5 □g/ml, Human Immunohistochemistry(Paraffin-embedded Section), 2-5 □g/ml, Human Immunocytochemistry/Immunofluorescence, 5 □g/ml, Human Flow Cytometry, 1-3 □g/1x^6 cells, Human
Source	Eukaryota
Contents	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na2HPO4.
Clone Names	Clone: 9H2F8
Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human Cytokeratin 5, different from the related mouse sequence by one amino acid, and identical to the related rat sequence.
Purification	Immunogen affinity purified.
Storage	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.

Protein Information

Name	KRT5 (HGNC:6442)
Function	Structural component of intermediate filaments in basal keratinocytes of stratified epithelia. Forms heteropolymers with type I partners including KRT14 and KRT15, contributing to the formation of the keratin intermediate filament network that provides mechanical stability and resilience to the basal layer of the epidermis (PubMed:1372711, PubMed:1694855). Regulates the recruitment of Langerhans cells to the epidermis, potentially by modulation of the abundance of macrophage chemotactic cytokines, macrophage inflammatory cytokines and CTNND1 localization in keratinocytes (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Expressed in corneal epithelium (at protein level) (PubMed:26758872). Expressed in keratinocytes (at protein level) (PubMed:20128788, PubMed:31302245).

Background

Cytokeratin 5, also known as KRT5, K5, or CK5, is a protein that is encoded in humans by the KRT5 gene. The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is specifically expressed in the basal layer of the epidermis with family member KRT14. Mutations in these genes have been associated with a complex of diseases termed epidermolysis bullosa simplex. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.

Images




Figure 1. Western blot analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7).
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.
Lane 1: human A431 whole cell lysates,
Lane 2: human SH-SY5Y whole cell lysates,
Lane 3: human Hacat whole cell lysates.
After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Cytokeratin 5 antigen affinity purified monoclonal antibody (Catalog # M00398-7) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Cytokeratin 5 at approximately 62 kDa. The expected band size for Cytokeratin 5 is at 62 kDa.




Figure 2. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7).
Cytokeratin 5 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 3. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7).
Cytokeratin 5 was detected in a paraffin-embedded section of human metaplasia of squamous cells of the renal pelvis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 4. IHC analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7).
Cytokeratin 5 was detected in a paraffin-embedded section of human rectal moderately differentiated adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 µg/ml mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.




Figure 5. IF analysis of Cytokeratin 5 using anti-Cytokeratin 5 antibody (M00398-7).
Cytokeratin 5 was detected in an immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 µg/mL mouse anti-Cytokeratin 5 Antibody (M00398-7) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. Flow Cytometry analysis of A431 cells using anti-Cytokeratin 5 antibody (M00398-7).
Overlay histogram showing A431 cells stained with M00398-7 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Cytokeratin 5 Antibody (M00398-7, 1 µg/1x10^6 cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 µg/1x10^6 cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 µg/1x10^6) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.