

Anti-Akt(pan) 1/2/3 Rabbit Monoclonal Antibody

Catalog # ABO15145

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	P31749/P31751/Q9Y243
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Akt(pan) 1/2/3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.


Additional Information

Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 17A20
Immunogen	A synthesized peptide derived from human AKT1/2/3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.


Protein Information

Images


All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:5K dilution for 1 hour at room temperature.




Western blot analysis of Akt(pan)1/2/3 expression in (1) A549 cell lysate; (2) Mouse brain lysate; (3) Rat brain lysate; (4) Rat heart lysate; (5) Rat kidney lysate.





Immunohistochemical analysis of paraffin-embedded Human pituitary tumor, using the Antibody at 1:250 dilution.




Immunohistochemical analysis of paraffin-embedded Human astrocytoma, using the Antibody at 1:250 dilution.




Immunohistochemical analysis of paraffin-embedded Human ovarian cancer, using the Antibody at 1:1000 dilution.



Immunohistochemical analysis of paraffin-embedded human testis, using Akt(pan)1/2/3 Antibody.



Immunofluorescent analysis of MCF-7 cells, using Akt(pan)1/2/3 Antibody.



Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:3K dilution)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.