

Anti-Factor X Rabbit Monoclonal Antibody

Catalog # ABO15228

Product Information

Application	WB, IHC
Primary Accession	P00742
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-Factor X Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human.

Additional Information

Gene ID	2159
Other Names	Coagulation factor X, 3.4.21.6, Stuart factor, Stuart-Prower factor, Factor X light chain, Factor X heavy chain, Activated factor Xa heavy chain, F10
Calculated MW	54732
Application Details	WB 1:1000-1:5000 IHC 1:1000-1:500
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18F04
Immunogen	A synthesized peptide derived from human Factor X
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	F10
Function	Factor Xa is a vitamin K-dependent glycoprotein that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting (PubMed: 22409427 , PubMed: 39880037). Factor Xa activates pro-inflammatory signaling pathways in a protease- activated receptor

(PAR)-dependent manner (PubMed:[24041930](#), PubMed:[30568593](#), PubMed:[34831181](#), PubMed:[18202198](#)). Up-regulates expression of protease-activated receptors (PARs) F2R, F2RL1 and F2RL2 in dermal microvascular endothelial cells (PubMed:[35738824](#)). Triggers the production of pro-inflammatory cytokines, such as MCP-1/CCL2 and IL6, in cardiac fibroblasts and umbilical vein endothelial cells in PAR-1/F2R-dependent manner (PubMed:[30568593](#), PubMed:[34831181](#)). Triggers the production of pro-inflammatory cytokines, such as MCP-1/CCL2, IL6, TNF/TNF, IL-1beta/IL1B, IL8/CXCL8 and IL18, in endothelial cells and atrial tissues (PubMed:[24041930](#), PubMed:[35738824](#), PubMed:[9780208](#)). Induces expression of adhesion molecules, such as ICAM1, VCAM1 and SELE, in endothelial cells and atrial tissues (PubMed:[24041930](#), PubMed:[35738824](#), PubMed:[9780208](#)). Increases expression of phosphorylated ERK1/2 in dermal microvascular endothelial cells and atrial tissues (PubMed:[24041930](#), PubMed:[35738824](#)). Triggers activation of the transcription factor NF-kappa-B in dermal microvascular endothelial cells and atrial tissues (PubMed:[24041930](#), PubMed:[35738824](#)). Activates pro-inflammatory and pro-fibrotic responses in dermal fibroblasts and enhances wound healing probably via PAR- 2/F2RL1-dependent mechanism (PubMed:[18202198](#)). Activates barrier protective signaling responses in endothelial cells in PAR-2/F2RL1- dependent manner; the activity depends on the cleavage of PAR-2/F2RL1 by factor Xa (PubMed:[22409427](#)). Up-regulates expression of plasminogen activator inhibitor 1 (SERPINE1) in atrial tissues (PubMed:[24041930](#)).


Cellular Location

Secreted.

Tissue Location

Plasma; synthesized in the liver.

Images



Western blot analysis of Factor X expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.