10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Niemann Pick C1 Rabbit Monoclonal Antibody

Catalog # ABO15231

## Product Information

| | |
|---|---|
| **Application** | WB, IHC, IF, ICC, FC |
| **Primary Accession** | O15118 |
| **Host** | Rabbit |
| **Isotype** | IgG |
| **Reactivity** | Rat, Human, Mouse |
| **Clonality** | Monoclonal |
| **Format** | Liquid |
| **Description** | Anti-Niemann Pick C1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. |

## Additional Information

| | |
|---|---|
| **Gene ID** | 4864 |
| **Other Names** | NPC intracellular cholesterol transporter 1 {ECO:0000312\|HGNC:HGNC:7897}, Niemann-Pick C1 protein, NPC1 (HGNC:7897) |
| **Calculated MW** | 142167 |
| **Application Details** | WB 1:500-1:2000<br>IHC 1:50-1:200<br>ICC/IF 1:50-1:200<br>FC 1:100 |
| **Source** | Eukaryota |
| **Contents** | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA. |
| **Clone Names** | Clone: 18N07 |
| **Immunogen** | A synthesized peptide derived from human Niemann Pick C1 |
| **Purification** | Affinity-chromatography |
| **Storage** | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles. |

## Protein Information


| | |
|---|---|
| **Name** | NPC1 ( HGNC:7897) |
| **Function** | Intracellular cholesterol transporter which acts in concert with NPC2 and plays an important role in the egress of cholesterol from the |

endosomal/lysosomal compartment (PubMed:10821832, PubMed:12554680, PubMed:18772377, PubMed:27238017, PubMed:9211849, PubMed:9927649). Unesterified cholesterol that has been released from LDLs in the lumen of the late endosomes/lysosomes is transferred by NPC2 to the cholesterol-binding pocket in the N-terminal domain of NPC1 (PubMed:18772377, PubMed:19563754, PubMed:27238017, PubMed:27378690, PubMed:28784760, PubMed:9211849, PubMed:9927649). Cholesterol binds to NPC1 with the hydroxyl group buried in the binding pocket (PubMed:19563754). Mediates cholesterol transfer from lysosomes to endoplasmic reticulum by tethering endosomal/lysosomal compartments and endoplasmic reticulum membrane contact sites where it interacts with sterol transport protein GRAMD1B (PubMed:31537798). Binds oxysterol with higher affinity than cholesterol. May play a role in vesicular trafficking in glia, a process that may be crucial for maintaining the structural and functional integrity of nerve terminals (Probable). Inhibits cholesterol-mediated mTORC1 activation throught its interaction with SLC38A9 (PubMed:28336668).

**Cellular Location**

Late endosome membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein

## Images




Western blot analysis of Niemann Pick C1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.