

Anti-C1QB Rabbit Monoclonal Antibody

Catalog # ABO15239

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P02746
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-C1QB Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information

Gene ID	713
Other Names	Complement C1q subcomponent subunit B, C1QB
Calculated MW	26722
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18C15
Immunogen	A synthesized peptide derived from human C1QB
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	C1QB {ECO:0000303 PubMed:3000358, ECO:0000312 HGNC:HGNC:1242}
Function	Core component of the complement C1 complex, a multiprotein complex that initiates the classical pathway of the complement system, a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed: 12847249 ,

PubMed:[19006321](#), PubMed:[24626930](#), PubMed:[29449492](#), PubMed:[3258649](#), PubMed:[34155115](#), PubMed:[6249812](#), PubMed:[6776418](#)). The classical complement pathway is initiated by the C1Q subcomplex of the C1 complex, which specifically binds IgG or IgM immunoglobulins complexed with antigens, forming antigen-antibody complexes on the surface of pathogens: C1QA, together with C1QB and C1QC, specifically recognizes and binds the Fc regions of IgG or IgM via its C1q domain (PubMed:[12847249](#), PubMed:[19006321](#), PubMed:[24626930](#), PubMed:[29449492](#), PubMed:[3258649](#), PubMed:[6776418](#)). Immunoglobulin-binding activates the proenzyme C1R, which cleaves C1S, initiating the proteolytic cascade of the complement system (PubMed:[29449492](#)). The C1Q subcomplex is activated by a hexamer of IgG complexed with antigens, while it is activated by a pentameric IgM (PubMed:[19706439](#), PubMed:[24626930](#), PubMed:[29449492](#)). The C1Q subcomplex also recognizes and binds phosphatidylserine exposed on the surface of cells undergoing programmed cell death, possibly promoting activation of the complement system (PubMed:[18250442](#)).

Cellular Location

Secreted. Cell surface. Note=Specifically binds IgG or IgM immunoglobulins complexed with antigens, forming antigen-antibody complexes on the surface of pathogens.

Images



Western blot analysis of C1QB expression in Human plasma lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.