

Anti-EB3 Rabbit Monoclonal Antibody

Catalog # ABO15280

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q9UPY8
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-EB3 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	22924
Other Names	Microtubule-associated protein RP/EB family member 3, EB1 protein family member 3, EBF3, End-binding protein 3, EB3, RP3, MAPRE3
Calculated MW	31982
Application Details	WB 1:1000-1:5000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 18M56 A synthesized peptide derived from human EB3
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	MAPRE3
Function	Plus-end tracking protein (+TIP) that binds to the plus-end of microtubules and regulates the dynamics of the microtubule cytoskeleton

(PubMed:[19255245](#), PubMed:[28814570](#)). Promotes microtubule growth (PubMed:[19255245](#), PubMed:[28814570](#)). May be involved in spindle function by stabilizing microtubules and anchoring them at centrosomes (PubMed:[19255245](#), PubMed:[28814570](#)). Also acts as a regulator of minus-end microtubule organization: interacts with the complex formed by AKAP9 and PDE4DIP, leading to recruit CAMSAP2 to the Golgi apparatus, thereby tethering non-centrosomal minus-end microtubules to the Golgi, an important step for polarized cell movement (PubMed:[28814570](#)). Promotes elongation of CAMSAP2-decorated microtubule stretches on the minus-end of microtubules (PubMed:[28814570](#)).


Cellular Location

Cytoplasm, cytoskeleton. Note=Associated with the microtubule network. Detected at the plus end of microtubules

Tissue Location

Predominantly expressed in brain and muscle.

Images



Western blot analysis of EB3 expression in Rat muscle cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.