

Anti-CYP26A1 Rabbit Monoclonal Antibody

Catalog # ABO15283

Product Information

Application	WB, IHC
Primary Accession	O43174
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-CYP26A1 Rabbit Monoclonal Antibody . Tested in WB, IHC applications. This antibody reacts with Human, Rat.

Additional Information

Gene ID	1592
Other Names	Cytochrome P450 26A1, CYP26A1, 1.14.13.-, Cytochrome P450 retinoic acid-inactivating 1, Cytochrome P450RAI, hP450RAI, Retinoic acid 4-hydroxylase, Retinoic acid-metabolizing cytochrome, CYP26A1 {ECO:0000303 PubMed:26937021, ECO:0000312 HGNC:HGNC:2603}
Calculated MW	56199
Application Details	WB 1:500-1:2000 IHC 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18C59
Immunogen	A synthesized peptide derived from human CYP26A1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CYP26A1 {ECO:0000303 PubMed:26937021, ECO:0000312 HGNC:HGNC:2603}
Function	A cytochrome P450 monooxygenase involved in the metabolism of

retinoates (RAs), the active metabolites of vitamin A, and critical signaling molecules in animals (PubMed:[22020119](#), PubMed:[9228017](#), PubMed:[9716180](#)). RAs exist as at least four different isomers: all- trans-RA (atRA), 9-cis-RA, 13-cis-RA, and 9,13-dicis-RA, where atRA is considered to be the biologically active isomer, although 9-cis-RA and 13-cis-RA also have activity (Probable). Catalyzes the hydroxylation of atRA primarily at C-4 and C-18, thereby contributing to the regulation of atRA homeostasis and signaling (PubMed:[22020119](#), PubMed:[9228017](#), PubMed:[9716180](#)). Hydroxylation of atRA limits its biological activity and initiates a degradative process leading to its eventual elimination (Probable). Involved in the conversion of atRA to all-trans-4-oxo-RA. Able to metabolize other RAs such as 9-cis, 13-cis and 9,13-di-cis RA (By similarity) (PubMed:[9228017](#)). Can oxidize all-trans-13,14- dihydroretinoate (DRA) to metabolites which could include all-trans-4- oxo-DRA, all-trans-4-hydroxy-DRA, all-trans-5,8-epoxy-DRA, and all-trans-18-hydroxy-DRA (By similarity). May play a role in the oxidative metabolism of xenobiotics such as tazarotenic acid (PubMed:[26937021](#)).


Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Tissue Location

Expressed in most fetal and adult tissues with highest levels in adult liver, heart, pituitary gland, adrenal gland, placenta and regions of the brain (PubMed:[9826557](#)). Expressed at high levels in lung, pancreas, skin and uterus (at protein level) (PubMed:[22020119](#)). Lower expression level is detected in spleen, kidney, intestine and adipose tissue (at protein level) (PubMed:[22020119](#)).

Images



Western blot analysis of CYP26A1 expression in HepG2 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.