

Anti-SF3B1 Rabbit Monoclonal Antibody

Catalog # ABO15284

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	O75533
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-SF3B1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	23451
Other Names	Splicing factor 3B subunit 1, Pre-mRNA-splicing factor SF3b 155 kDa subunit, SF3b155 {ECO:0000303 Ref.39}, Spliceosome-associated protein 155, SAP 155, SF3B1 {ECO:0000303 PubMed:30567737, ECO:0000312 HGNC:HGNC:10768}
Calculated MW	145830
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:500
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18S60
Immunogen	A synthesized peptide derived from human SF3B1
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SF3B1 {ECO:0000303 PubMed:30567737, ECO:0000312 HGNC:HGNC:10768}
-------------	--


Function

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:[12234937](#), PubMed:[27720643](#), PubMed:[32494006](#), PubMed:[34822310](#), PubMed:[36104565](#)). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:[32494006](#), PubMed:[34822310](#), PubMed:[36104565](#)). Within the 17S U2 SnRNP complex, SF3B1 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed:[12234937](#), PubMed:[36104565](#)). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:[12234937](#)). May also be involved in the assembly of the 'E' complex (PubMed:[10882114](#)). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed:[15146077](#), PubMed:[33509932](#)). Together with other U2 snRNP complex components may also play a role in the selective processing of microRNAs (miRNAs) from the long primary miRNA transcript, pri-miR-17-92 (By similarity).

Cellular Location

Nucleus. Nucleus speckle. Note=During mitosis, transiently dispersed from the nuclear speckles to the cytoplasm

Images



Western blot analysis of SF3B1 expression in Jurkat cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.