

Anti-SLC4A1/CD233/Band 3 Rabbit Monoclonal Antibody

Catalog # ABO15317

Product Information

Application	WB, IHC, IF, ICC
Primary Accession	P02730
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-SLC4A1 Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.

Additional Information


Gene ID	6521
Other Names	Band 3 anion transport protein, Anion exchange protein 1, AE 1, Anion exchanger 1, Solute carrier family 4 member 1, CD233, SLC4A1 (HGNC:11027), AE1, DI, EPB3
Calculated MW	101792
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18S93
Immunogen	A synthesized peptide derived from human Band 3 / CD233
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SLC4A1 (HGNC:11027)
Synonyms	AE1, DI, EPB3

Function	Functions both as a transporter that mediates electroneutral anion exchange across the cell membrane and as a structural protein (PubMed:10926824, PubMed:14734552, PubMed:1538405, PubMed:16227998, PubMed:20151848, PubMed:24121512, PubMed:28387307, PubMed:35835865). Component of the ankyrin-1 complex of the erythrocyte membrane; required for normal flexibility and stability of the erythrocyte membrane and for normal erythrocyte shape via the interactions of its cytoplasmic domain with cytoskeletal proteins, glycolytic enzymes, and hemoglobin (PubMed:1538405, PubMed:20151848, PubMed:35835865). Functions as a transporter that mediates the 1:1 exchange of inorganic anions across the erythrocyte membrane. Mediates chloride-bicarbonate exchange in the kidney, and is required for normal acidification of the urine (PubMed:10926824, PubMed:14734552, PubMed:16227998, PubMed:24121512, PubMed:28387307).
Cellular Location	Cell membrane; Multi-pass membrane protein. Basolateral cell membrane; Multi-pass membrane protein. Note=Detected in the erythrocyte cell membrane and on the basolateral membrane of alpha-intercalated cells in the collecting duct in the kidney.
Tissue Location	Detected in erythrocytes (at protein level).

Images



Western blot analysis of Band 3 / CD233 expression in Red blood cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.