

Anti-CHD1L Rabbit Monoclonal Antibody

Catalog # ABO15320

Product Information

Application	WB, IHC, IF, ICC, IP, FC
Primary Accession	Q86WJ1
Host	Rabbit
Isotype	IgG
Reactivity	Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-CHD1L Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.

Additional Information


Gene ID	9557
Other Names	Chromodomain-helicase-DNA-binding protein 1-like, 3.6.4.-, Amplified in liver cancer protein 1, CHD1L {ECO:0000303 PubMed:34210977, ECO:0000312 HGNC:HGNC:1916}
Calculated MW	101000
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:40 FC 1:200
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 18C96
Immunogen	A synthesized peptide derived from human CHD1L
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	CHD1L {ECO:0000303 PubMed:34210977, ECO:0000312 HGNC:HGNC:1916}
-------------	---

Function	ATP-dependent chromatin remodeler that mediates chromatin- remodeling following DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 33357431 , PubMed: 34210977 , PubMed: 34486521 , PubMed: 34874266). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653 , PubMed: 34486521 , PubMed: 34874266). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed: 19661379 , PubMed: 29220652 , PubMed: 29220653). Chromatin remodeling activity promotes PARP2 removal from chromatin (PubMed: 33275888).
Cellular Location	Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.
Tissue Location	Frequently overexpressed in hepatomacellular carcinomas.

Images



Western blot analysis of CHD1L expression in A549 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.