

Anti-KMT1B / SUV39H2 Rabbit Monoclonal Antibody

Catalog # ABO15355

Product Information

Application	WB, IHC, IP
Primary Accession	Q9H5I1
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-KMT1B / SUV39H2 Rabbit Monoclonal Antibody . Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	79723
Other Names	Histone-lysine N-methyltransferase SUV39H2, 2.1.1.355, Histone H3-K9 methyltransferase 2, H3-K9-HMTase 2, Lysine N-methyltransferase 1B, Suppressor of variegation 3-9 homolog 2, Su(var)3-9 homolog 2, SUV39H2, KMT1B
Calculated MW	46682
Application Details	WB 1:500-1:1000 IHC 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19S31
Immunogen	A synthesized peptide derived from human KMT1B / SUV39H2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SUV39H2
Synonyms	KMT1B


Function

Histone methyltransferase that specifically mediates trimethylation of 'Lys-9' of histone H3 (H3K9me3) using monomethylated H3 'Lys-9' (H3K9me1) as substrate (PubMed:[10949293](#)). H3 'Lys-9' trimethylation represents a specific tag for epigenetic transcriptional repression by recruiting HP1 (CBX1, CBX3 and/or CBX5) proteins to methylated histones (PubMed:[10949293](#)). Mainly functions in heterochromatin regions, thereby playing a central role in the establishment of constitutive heterochromatin at pericentric and telomere regions (PubMed:[10949293](#)). H3 'Lys-9' trimethylation is also required to direct DNA methylation at pericentric repeats (PubMed:[10949293](#), PubMed:[41094145](#)). SUV39H2 is targeted to histone H3 via its interaction with RB1 and is involved in many processes, such as cell cycle regulation, transcriptional repression and regulation of telomere length (PubMed:[14765126](#)). May participate in regulation of higher-order chromatin organization during spermatogenesis (By similarity). Recruited by the large PER complex to the E-box elements of the circadian target genes such as PER2 itself or PER1, contributes to the conversion of local chromatin to a heterochromatin-like repressive state through H3 'Lys-9' trimethylation (By similarity).

Cellular Location

Nucleus. Chromosome. Chromosome, centromere {ECO:0000250|UniProtKB:Q9EQQ0}. Note=Associates with centromeric constitutive heterochromatin (By similarity). Recruited to pericentromeric heterochromatin via its chromo domain, which specifically recognizes and binds histone H3 monoubiquitinated at 'Lys- 14' (H3K14ub) (PubMed:41094145). {ECO:0000250|UniProtKB:Q9EQQ0, ECO:0000269|PubMed:41094145}

Images



Western blot analysis of KMT1B / SUV39H2 expression in (1) MOLT-4 cell lysate; (2) Human testis lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.