

Anti-SOCS2 Rabbit Monoclonal Antibody

Catalog # ABO15359

Product Information

Application	WB, IF, ICC, IP
Primary Accession	O14508
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-SOCS2 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP applications. This antibody reacts with Human.

Additional Information

Gene ID	8835
Other Names	Suppressor of cytokine signaling 2, SOCS-2, Cytokine-inducible SH2 protein 2, CIS-2, STAT-induced STAT inhibitor 2, SSI-2, SOCS2, CIS2, SSI2, STATI2
Calculated MW	22172
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 19S35 A synthesized peptide derived from human SOCS2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	SOCS2 {ECO:0000303 PubMed:10512686, ECO:0000312 HGNC:HGNC:19382}
Function	Substrate-recognition component of a cullin-5-RING E3 ubiquitin-protein ligase complex (ECS complex, also named CRL5 complex), which mediates the

ubiquitination and subsequent proteasomal degradation of target proteins, such as EPOR and GHR (PubMed:[11781573](#), PubMed:[21980433](#), PubMed:[25505247](#), PubMed:[31182716](#), PubMed:[34857742](#)). Specifically recognizes and binds phosphorylated proteins via its SH2 domain, promoting their ubiquitination (PubMed:[21980433](#), PubMed:[25505247](#), PubMed:[31182716](#), PubMed:[34857742](#), PubMed:[37816714](#)). The ECS(SOCS2) complex acts as a key regulator of growth hormone receptor (GHR) levels by mediating ubiquitination and degradation of GHR, following GHR phosphorylation by JAK2 (PubMed:[21980433](#), PubMed:[25505247](#), PubMed:[34857742](#)). The ECS(SOCS2) also catalyzes ubiquitination and degradation of JAK2-phosphorylated EPOR (PubMed:[11781573](#)).


Cellular Location

Cytoplasm.


Tissue Location

High expression in heart, placenta, lung, kidney and prostate. Predominantly expressed in pulmonary epithelia cells, specifically type II pneumocytes.

Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of SOCS2 expression in K562 cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.