

Anti-RHEB Rabbit Monoclonal Antibody

Catalog # ABO15364

Product Information

Application	WB, IHC, IF, ICC, FC
Primary Accession	Q15382
Host	Rabbit
Isotype	IgG
Reactivity	Human
Clonality	Monoclonal
Format	Liquid
Description	Anti-RHEB Rabbit Monoclonal Antibody . Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.

Additional Information

Gene ID	6009
Other Names	GTP-binding protein Rheb, Ras homolog enriched in brain, 3.6.5.-, RHEB {ECO:0000303 PubMed:8543055, ECO:0000312 HGNC:HGNC:10011}
Calculated MW	20497
Application Details	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 FC 1:100
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names	Clone: 19R40
Immunogen	A synthesized peptide derived from human RHEB
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information

Name	RHEB {ECO:0000303 PubMed:8543055, ECO:0000312 HGNC:HGNC:10011}
Function	Small GTPase that acts as an allosteric activator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates

anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:[12172553](#), PubMed:[12271141](#), PubMed:[12842888](#), PubMed:[12869586](#), PubMed:[12906785](#), PubMed:[15340059](#), PubMed:[15854902](#), PubMed:[16098514](#), PubMed:[20381137](#), PubMed:[22819219](#), PubMed:[24529379](#), PubMed:[29416044](#), PubMed:[32470140](#), PubMed:[33157014](#), PubMed:[25816988](#)). In response to nutrients, growth factors or amino acids, specifically activates the protein kinase activity of MTOR, the catalytic component of the mTORC1 complex: acts by causing a conformational change that allows the alignment of residues in the active site of MTOR, thereby enhancing the phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) (PubMed:[29236692](#), PubMed:[33157014](#)). RHEB is also required for localization of the TSC-TBC complex to lysosomal membranes (PubMed:[24529379](#)). In response to starvation, RHEB is inactivated by the TSC-TBC complex, preventing activation of mTORC1 (PubMed:[24529379](#), PubMed:[33157014](#)). Has low intrinsic GTPase activity (PubMed:[15340059](#)).


Cellular Location

Endomembrane system; Lipid-anchor; Cytoplasmic side. Lysosome membrane; Lipid-anchor; Cytoplasmic side. Golgi apparatus membrane; Lipid-anchor; Cytoplasmic side. Endoplasmic reticulum membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytosol. Note=Farnesylation is required for recruitment to lysosomal membranes, where it activates the mTORC1 complex.

Tissue Location

Ubiquitous (PubMed:[8543055](#)). Highest levels observed in skeletal and cardiac muscle (PubMed:[8543055](#))

Images



Western blot analysis of RHEB expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.