

Anti-Cullin 2 Rabbit Monoclonal Antibody

Catalog # ABO15367

Product Information

Application	WB, IF, ICC, IP, FC
Primary Accession	Q13617
Host	Rabbit
Isotype	IgG
Reactivity	Rat, Human, Mouse
Clonality	Monoclonal
Format	Liquid
Description	Anti-Cullin 2 Rabbit Monoclonal Antibody . Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.

Additional Information

Gene ID	8453
Other Names	Cullin-2, CUL-2, CUL2 (HGNC:2552)
Calculated MW	86983
Application Details	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
Source	Eukaryota
Contents	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Clone Names Immunogen	Clone: 19C43 A synthesized peptide derived from human Cullin 2
Purification	Affinity-chromatography
Storage	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.

Protein Information


Name	CUL2 (HGNC:2552)
Function	Core component of multiple cullin-RING-based ECS (ElonginB/C-CUL2/5-SOCS-box protein) E3 ubiquitin-protein ligase complexes, which mediate the ubiquitination of target proteins (PubMed: 11384984 ,

PubMed:[26138980](#), PubMed:[29775578](#), PubMed:[29779948](#), PubMed:[37844242](#), PubMed:[38326650](#)). CUL2 serves as a rigid scaffold in the complex and may contribute to catalysis through positioning of the substrate and the E2 ubiquitin-conjugating enzyme (PubMed:[10973499](#), PubMed:[11384984](#), PubMed:[12609982](#), PubMed:[24076655](#), PubMed:[9122164](#), PubMed:[37844242](#), PubMed:[38326650](#)). The E3 ubiquitin-protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1 (PubMed:[12609982](#), PubMed:[24076655](#), PubMed:[27565346](#), PubMed:[38326650](#)). The functional specificity of the ECS complex depends on the substrate recognition component (PubMed:[10973499](#), PubMed:[26138980](#), PubMed:[29775578](#), PubMed:[29779948](#), PubMed:[9122164](#), PubMed:[38326650](#)). ECS(VHL) mediates the ubiquitination of hypoxia-inducible factor (HIF) (PubMed:[10973499](#), PubMed:[9122164](#)). A number of ECS complexes (containing either KLHDC2, KLHDC3, KLHDC10, APPBP2, FEM1A, FEM1B or FEM1C as substrate-recognition component) are part of the DescEND (destruction via C-end degrons) pathway, which recognizes a C-degron located at the extreme C terminus of target proteins, leading to their ubiquitination and degradation (PubMed:[26138980](#), PubMed:[29775578](#), PubMed:[29779948](#), PubMed:[37844242](#)). ECS complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed:[27565346](#)). ECS(LRR1) ubiquitinates MCM7 and promotes CMG replisome disassembly by VCP and chromatin extraction during S- phase (By similarity).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9D4H8}.

Images



Western blot analysis of Cullin 2 expression in Raji cell lysate.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.